



Leibniz Institute
for Prevention Research and
Epidemiology – BIPS

Dietary fat, the gut microbiota, and metabolic health - A systematic review conducted within the MyNewGut project

Maike Wolters, Jenny Ahrens, Marina Romaní-Perez, Claire Watkins, Yolanda Sanz, Alfonso Benítez-Paez, Cathrine Stanton, Kathrin Günther

DOI

10.1016/j.clnu.2018.12.024

Published in

Clinical Nutrition

Document version

Accepted manuscript

This is the author's final accepted version. There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

Online publication date

24 December 2018

Corresponding author

Maike Wolters

Citation

Wolters M, Ahrens J, Romaní-Perez M, Watkins C, Sanz Y, Benítez-Paez A, et al. Dietary fat, the gut microbiota, and metabolic health - A systematic review conducted within the MyNewGut project. Clin Nutr. 2019;38(6):2504-20.



© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

1 **Dietary fat, the gut microbiota, and metabolic health – a systematic review conducted within the**
2 **MyNewGut project**

3 Maïke Wolters ^{a#}, Jenny Ahrens ^a, Marina Romani Perez ^b, Claire Watkins ^c, Yolanda Sanz ^b, Alfonso
4 Benítez-Páez ^b, Catherine Stanton ^{c*}, Kathrin Günther ^{a*}

5

6 ^a Leibniz Institute for Prevention Research and Epidemiology – BIPS, Achterstraße 30, 28359 Bremen,
7 Germany

8 ^b Microbial Ecology, Nutrition & Health Research Unit. Institute of Agrochemistry and Food
9 Technology, National Research Council (IATA-CSIC), Valencia, Spain

10 ^c APC Microbiome Institute, University College Cork, Cork, Ireland

11

12 **#Corresponding author:**

13 Dr. rer. nat. Maïke Wolters

14 Leibniz Institute for Prevention Research and Epidemiology - BIPS

15 Department Epidemiological Methods and Etiological Research

16 Achterstr. 30

17 28359 Bremen

18 Germany

19 Tel: +49 / 421 / 218-56845

20 Fax: +49 / 421 / 218-56821

21 E-Mail: wolters@leibniz-bips.de

22

23 *Shared senior authorship

24 Abbreviations: ACE, abundance-based coverage estimation; ALA, α -linolenic acid; BF%, body fat
25 percentage; BMI, body mass index; BP, blood pressure; CG, control group; CHD, coronary heart
26 disease; CHO, carbohydrates; CI, confidence interval; CV, cardiovascular; d, day; DGGE, Denaturing
27 Gradient Gel Electrophoresis; DHA, docosahexaenoic acid; EPA, eicosapentanoic acid; *EreC*,
28 *Eubacterium rectale-Clostridium coccooides*; E%, energy percentage; FBG, fasting blood glucose; F/B,
29 Firmicutes to Bacteroides; FISH, fluorescence in-situ hybridization; GI, glycemic index; HDL, high-
30 density lipoprotein cholesterol; HOMA, Homeostasis model of insulin resistance; IG, intervention
31 group; IQ, interquartile range; KO, Knockout; LA, linoleic acid; LDL, low-density lipoprotein
32 cholesterol; LFHCC diet, low-fat, high-complex carbohydrate diet; LN, lean; LOM, logarithmic orders
33 of magnitude; Med diet, Mediterranean diet; MetS, Metabolic Syndrome; MUFA, monounsaturated
34 fatty acids; NEFA, non-esterified fatty acids; NOS, Newcastle-Ottawa Scale; n3 PUFA, omega-3
35 polyunsaturated fatty acid; n6 PUFA, omega-6 polyunsaturated fatty acid; NW, normal weight; OB,
36 obese; OTU, operational taxonomic unit; OW, overweight; PCoA, Principal coordinate analysis; PICO,
37 Population-Intervention-Comparison-Outcome; PRISMA, Preferred Reporting Items for Systematic
38 reviews and Meta-Analysis; PCR, polymerase chain reaction; PUFA, polyunsaturated fatty acids;
39 qPCR, quantitative polymerase chain reaction; RCT, randomized controlled trial; resp., respectively;
40 SCFA, short-chain fatty acids; SE, standard error; SEM, standard error of the mean; SFA, saturated
41 fatty acid; TC, total cholesterol; TG, triglycerides; T2D, Type 2 diabetes; USFA, unsaturated fatty
42 acids; UW, underweight; VLDL, very low density lipoprotein; VO₂max, maximum oxygen uptake; WC,
43 waist circumference; y, year.

44

45 **Abstract**

46 **Background and aims:** Studies indicate that dietary fat quantity and quality influence the gut
47 microbiota composition which may as a consequence impact metabolic health. This systematic
48 review aims to summarize the results of available studies in humans on dietary fat intake (quantity
49 and quality), the intestinal microbiota composition and related cardiometabolic health outcomes.

50 **Methods:** We performed a systematic review (CRD42018088685) following PRISMA guidelines and
51 searched for literature in Medline, EMBASE, and Cochrane databases.

52 **Results:** From 796 records, 765 records were excluded based on title or abstract. After screening of
53 31 full-text articles six randomized controlled trials (RCT) and nine cross-sectional observational
54 studies were included. Our results of interventional trials do not suggest strong effects of different
55 amounts and types of dietary fat on the intestinal microbiota composition or on metabolic health
56 outcomes while observational studies indicate associations with the microbiota and health
57 outcomes. High intake of fat and saturated fatty acids (SFA) may negatively affect microbiota
58 richness and diversity and diets high in monounsaturated fatty acids (MUFA) may decrease total
59 bacterial numbers whereas dietary polyunsaturated fatty acids (PUFA) had no effect on richness and
60 diversity.

61 **Conclusions:** High fat and high SFA diets can exert unfavorable effects on the gut microbiota and are
62 associated with an unhealthy metabolic state. Also high MUFA diets may negatively affect gut
63 microbiota whereas PUFA do not seem to negatively affect the gut microbiota or metabolic health
64 outcomes. However, data are not consistent and most RCT and observational studies showed risks of
65 bias.

66

67 **Keywords**

68 Fat; fatty acids; saturated fatty acids; unsaturated fatty acids; intestinal microbiota; metabolic health

69

70 **Introduction and rationale**

71 In recent years, the gut microbiota has emerged as a significant factor for the regulation of energy
72 balance and has been shown to be associated with obesity and metabolic diseases. The gut
73 microbiota plays an important role in polysaccharide fermentation and the production of short-chain
74 fatty acids (SCFA) which can be metabolized or used for the *de novo* synthesis of glucose, lipids or
75 bile acids [1, 2]. Additionally the gut microbiota is involved in the maintenance of barrier function of
76 the intestinal epithelium preventing the translocation of lipopolysaccharides and related
77 endotoxemia which can lead to inflammation and increased risk of insulin resistance [3, 4]. Studies
78 indicated that impaired gut microbiota-host interactions at infancy, e.g. by antibiotic use could
79 increase the risk of metabolic diseases in later life [5].

80

81 *Dietary intervention and metabolic health outcome*

82 Dietary sources of energy and nutrients play a significant role in the development of obesity and
83 metabolic diseases and also modulate the gut microbiota. Theoretically, dietary-induced microbiota
84 changes could also be partly responsible for the metabolic phenotype of the person. Indeed, the
85 obese microbiome has previously been reported to have an increased capacity to harvest energy
86 from the diet when transferred from humans to germ-free mice [6]. In observational studies in
87 humans, *Bacteroides* spp., *Bilophila wadsworthia* and *Alistipes* have been associated with a long-term
88 diet high in animal protein and saturated fats, whereas *Prevotella*, *Roseburia*, *Eubacterium rectale*
89 and *Facalibacterium prausnitzii* have been associated with plant-based diets high in carbohydrates
90 and simple sugars [1, 7, 8]. Animal studies indicate that high fat diets are associated with changes in
91 the gut microbiota leading to inflammation and increased risk of insulin resistance. In particular,
92 high-fat diets rich in long-chain saturated fatty acids (SFA) have been found to modulate the gut
93 microbiota resulting in dysbiosis, inflammation and consequently an increased risk of obesity and
94 metabolic syndrome (MetS) [9, 10]. In contrast, beneficial effects were observed for high tissue levels
95 of n3 polyunsaturated fatty acids (PUFA) which reduced body weight gain and the severity of insulin
96 resistance, fatty liver and dyslipidemia resulting from early-life exposure to antibiotics in a mouse

97 model [11] but effects on microbiota are less well documented. Selective enrichment of specific
98 microorganisms has also been found to promote metabolic health in a number of dietary
99 intervention studies in humans [12, 13].

100

101 *The gut microbiome and metabolic biomarkers in at-risk populations*

102 An altered gut microbiome has been reported in individuals with type 2 diabetes, independent of
103 body mass index (BMI) [14, 15]. When compared to individuals with normal glucose tolerance, an
104 increased abundance of *Lactobacillus* spp. and a decreased abundance of *Clostridium* spp. were
105 shown in individuals with type 2 diabetes [14]. Furthermore, a mathematical model based on
106 shotgun metagenomic profiles identified an increase in *Clostridium clostridioforme* and a decrease in
107 *Roseburia* 272 metagenomic clusters in type 2 diabetes from two cohorts [14, 16]. Depletion of
108 *Akkermansia muciniphilia* has also been described as a microbial biomarker for type 2 diabetes prior
109 to the onset of disease in a metagenomics study in monozygotic Korean twins [15]. Reduced butyrate
110 and a decreased abundance of butyrate-producing genera, such as *Roseburia*, *Faecalibacterium* and
111 *Clostridium*, have been found to be associated with obesity and impaired glucose tolerance [17, 18].
112 Interestingly, an increase in propionate-producing genera, such as *Bacteroides* and *Prevotella*, were
113 found in overweight and obese human individuals [19], suggesting a potential inverse relationship
114 between butyrate and propionate with regard to metabolic health although also beneficial effects of
115 propionate on metabolic health have been reported [17, 20].

116 Thus this systematic literature review intends to investigate effects of dietary fat quantity and quality
117 including different types of fatty acids on the gut microbiota and metabolic health outcomes in
118 humans. It was performed within MyNewGut (<http://www.mynewgut.eu/>), a FP7 EU project which
119 aims to disentangle the role played by the gut microbiota (via interactions with lifestyle factors, e.g.
120 diet, eating habits, stress, etc.), in the regulation of pathways leading to the development of obesity
121 and the associated metabolic and behavioral disorders. This review is part of a series of position
122 papers of the MyNewGut project aiming at informing future recommendations for dietary guidelines

123 based on project results and the latest advantages in the field regarding insights gained in the role of
124 the gut microbiome.

125

126 **Methods**

127 *Search strategy and in-/exclusion criteria*

128 We performed a systematic literature review following Preferred Reporting Items for Systematic
129 reviews and Meta-Analysis (PRISMA) guidelines [21]. Our review protocol was registered on
130 PROSPERO under the Registration Number CRD42018088685.

131 To identify studies we searched for literature in Medline via PubMed, EMBASE via the Elsevier
132 platform, and the Cochrane databases via Wiley from their inception. All searches were performed
133 on January 17, 2018 using a combination of subject and free-text terms with no date limit or
134 language restriction. The search strategy was developed for Medline and adapted to yield results in
135 other databases. Details on the Medline search strategy are provided in Supplementary Table 1.

136 Eligibility criteria included dietary fat or fatty acids as exposure of interest, the composition of the
137 intestinal microbiota, and metabolic health markers such as MetS score, overweight/obesity,
138 increased waist circumference, insulin resistance, hypertension, dyslipidemia or cardiovascular
139 diseases as outcomes. After extraction of the references, the following four criteria were considered
140 for further evaluation of an abstract: a) an experimental or observational comparative study in
141 humans, b) diets varying in composition or quantity of fat or fatty acid intake including biomarkers
142 for the intake, e.g. serum level of PUFA, c) association with or effects on the gut microbiota
143 composition, d) a metabolic health outcome in terms of the MetS, any of its components or
144 cardiovascular diseases. Study exclusions were no study in humans, a review and/or meta-analysis,
145 insufficient information on the quantity and/or quality of dietary fat or fatty acids or on the gut
146 microbiota composition or on the metabolic outcome. Guidelines, editorials, case-reports,

147 dissertations or unpublished studies as well as conference abstracts and conference proceedings
148 were not considered.

149 Titles, abstracts and full-texts of articles were screened independently by two reviewers (MW, JA) for
150 eligibility. Disagreement was resolved by discussion and by a third senior reviewer (KG), when
151 needed.

152

153 *Data extraction*

154 The following data were extracted from each included study: first author's last name, publication
155 year, country, information on study design, number and characteristics of participants, dietary fat/
156 fatty acid intake and/or biomarkers, intestinal microbiota composition, weight status, metabolic
157 health outcomes, and follow-up time. Data extraction was performed independently by pairs of
158 reviewers (MW, JA). A third reviewer (KG) resolved disagreement if needed.

159

160 *Assessment of risk of bias*

161 Quality assessment of risk of bias of randomized controlled trials (RCT) was conducted using the
162 Cochrane risk of bias tool [22] through which selection, performance, detection, attrition and
163 reporting bias of each study were judged as high, low or unclear risk. The assessment of
164 observational studies was performed using the Newcastle-Ottawa Scale (NOS) [23] that evaluates 9
165 items grouped in 3 domains: selection of participants (maximum score 4 stars), comparability of
166 groups (maximum score 2 stars) and ascertainment of the outcomes of interest (maximum score 4
167 stars). Total score ranged from 0 to 9 and higher score indicated better methodological quality. Two
168 reviewers (MRP, CW) independently assessed the risk of bias of individual studies and any
169 differences in quality assessment results were resolved through consensus.

170

171 *Data synthesis and analysis*


172 As study designs and outcome assessments varied, results are presented in a narrative way. Studies
 173 are presented based on the PICO criteria (Population, Intervention [or Exposure], Comparison [if
 174 applicable], Outcome).

175

176

177 Results

178 Figure 1 shows the flow diagram of the screened and selected studies.



179

180 Figure 1: Flow chart of the selection process

181

182 Fifteen studies were included in this systematic review. Table 1 summarizes the characteristics and
 183 results of the interventional studies and Table 2 of the observational studies included. Six of the
 184 studies evaluated dietary interventions [24-29], five reviewed dietary records [30-35], one
 185 investigated serum metabolites of fatty acids [36] and three applied a food frequency questionnaire

186 [35, 37, 38]. All of the included studies were published between 2013 and 2017. Distinct study
187 designs were found, with six RCT [24-29], seven cross-sectional studies [30-32, 34, 35, 37, 38], and
188 two longitudinal cohort studies which analyzed cross-sectional data [33, 36]. Geographically, nine of
189 the studies were performed in Europe [24-27, 30, 36], three in North America [28, 31, 37] and three
190 in Asia [29, 35, 38]. Considering the patient selection, ten studies had no gender limit, three included
191 only women [30, 32, 34] and two included only men [27, 36]. Total sample sizes ranged from 20 [27]
192 to 88 [26] in interventional and from nine [37] to 531 [36] in observational studies. The mean age of
193 participants in the included studies varied between 8.1 and 63.3 years. The length of the
194 interventions varied from three weeks to one year, with a follow-up time of up to six months. With
195 one exception [28], in all interventional studies, a baseline assessment of the gut microbiota was
196 obtained, and microbial compositional changes were reported.

197

198 Table 1: Characteristics and results of the randomized controlled interventional studies included in the systematic review

Study	Population		Study design and intervention		Results / Effects on outcomes	
Reference	N	Characteristics	Description	Fat intake	Microbiota	Metabolic health
Balfegó et al. 2016 [24]	N=35 (32 finished) F: 54.3% IG: N=19 CG:N=16	T2D patients Mean (SE) Age: 60.6 (1.4) y BMI: IG 30.5 (1.0) kg/m ² CG 28.8 (0.8) kg/m ²	Randomized controlled nutritional pilot trial Country: Spain 2-week lead-in period, then 6 months dietary intervention; 1 visit/month by the dietician IG: standard diet for T2D enriched with 100 g of sardines (instead of usual protein foods) on 5 d/week CG: standard T2D diet	Daily intake, mean (SE), IG / CG: Fat (g): 88.3 (4.8) / 79.2 (4.5) (Baseline) 84.4 (8.1) / 83.7 (5.8) (6 months) PUFA (E%): 5.8 (0.4) / 7.1 (0.5) (Baseline) 6.2 (0.4) / 6.3 (0.4) (6 months) IG from sardines, g: n3 PUFA: 3.5 (0.2) EPA+DHA: 3.0 (0.2)	No differences in the abundance of the bacterial groups analyzed comparing IG and CG at 6 months Changes after 6 months compared to baseline: IG: ↓ <i>Firmicutes</i> ↑ <i>Escherichia coli</i> ↑ <i>Bacteroides-Prevotella</i> CG: ↓ <i>Firmicutes</i> ↑ <i>Escherichia coli</i> Trend of ↓ <i>Firmicutes/Bacteroidetes</i> ratio	Both groups: ↓ Fasting insulin and HOMA (compared to baseline) but mean change from baseline to 6 months was not different between IG and CG - IG patients exhibited greater decrease from baseline: IG: -6.1±1.8 mU/l insulin, -2.3±0.7 HOMA CG: -3.4±1.5 mU/l insulin, -1.1±0.7 HOMA CG: ↓ HbA1c (-0.3±0.1%) IG: non-significant ↓HbA1c (-0.2±0.1%)
Blædel et al. 2016 [25]	N=21 (18 finished) F: 0	Healthy men, aged 23-45 y Mean (SE) Age: 32.9 (0.85) y BMI: 29.3 (0.5) kg/m ²	Randomized, controlled, crossover study Country: Denmark 21 d intervention periods, separated by a wash-out period 3 arms: Isoenergetic standard diet with either - whole-fat milk (IG) - water (CG) - inulin powder (not considered here)	IG / CG (E%): CHO: 45 / 55 Fat: 40 / 35 Protein: 15 / 15	The overall fecal microbiota composition did not change significantly in response to milk (IG) compared with CG.	- No change in blood lipid profile, insulin or glucose concentration in IG compared to CG - No effect of diets on resting energy expenditure and lipid oxidation.
Fava et al. 2013 [26]	N=88 F: 51.1%	Adults at increased risk for MetS Mean (SD) Age: 54.0 (9.5) y	Randomized, controlled, single blind, parallel design Country: United	High fat diets: IG1 / IG2 / CG (E%): Total fat: 38 / 38 / 38 SFA: 10 / 10 / 18	↓ Total bacteria after intervention in the 3 diets with highest fat content (CG, IG1, IG2) compared to baseline ↓ Total bacterial numbers after both high MUFA-	- No significant changes in BMI, WC, BF% or BP between the diets at the end of intervention - No effect of the dietary interventions on insulin sensitivity parameters

		BMI: 28.8 (4.9) kg/m ² HDL: 1.6 (0.4) mmol/l	Kingdom 4-week run-in reference diet (CG, baseline: after run-in), then 24 weeks of one of the diets (matched for age, BMI, HDL) CG: reference diet IG1: HM/HGI: high MUFA/high GI IG2: HM/LGI: high MUFA/low GI IG3: HC/HGI: high CHO/high GI IG4: HC/LGI: high CHO/low GI	MUFA: 20 / 20 / 12 PUFA: 6 / 6 / 6 CHO: 45 / 45 / 45 GI: 64 / 53 / 64 High CHO diets: IG3 / IG4 (E%): Total fat: 28 / 28 SFA: 10 / 10 MUFA: 11 / 11 PUFA: 6 / 6 CHO: 55 / 55 GI: 64 / 51	diets (IG1, IG2) compared with IG3 and with baseline ↑ <i>Faecalibacterium prausnitzii</i> after intervention with CG compared to baseline and IG4 ↑ <i>Bifidobacterium</i> spp. population levels in IG3 compared to CG ↑ <i>Bifidobacterium</i> spp. population levels in IG3 and IG4 diets compared to baseline ↑ <i>Bacteroides</i> spp. in IG3 compared to baseline, but not compared to the other diets	↓ in NEFA concentration after intervention with IG4 compared to CG and to IG3 After treatment compared to baseline: ↓ in WC in IG2 ↓ in TC and LDL in all intervention groups ↓ in BF% in IG3 ↓ in HDL after IG3 ↓ FBG after IG3 and IG4 ↓ Plasma insulin after IG3 ↑ NEFA after IG3 ↓ NEFA after IG4 ↑ <i>Bacteroides</i> spp. numbers after IG3 diet was associated with decreases in body weight, BMI and WC (r=-0.64, r=-0.64 and r=- 0.45, resp.)
Haro et al. 2016 [27]	N=20 F: 0	Obese CHD patients Mean (SE) Age: 63.3 (2.0) y BMI: 32.2 (0.5) kg/m ²	Interventional study Country: Spain Participants received either a low-fat, high-complex CHO diet (LFHCC) or a Mediterranean diet (Med diet) for 1 year	LFHCC diet / Med diet (E%): Fat: 28 / 35 MUFA: 12 / 22 PUFA: 8 / 6 SFA: 8 / 7	LFHCC diet compared to baseline: ↑ <i>Prevotella</i> ↓ <i>Roseburia</i> ↑ <i>Faecalibacterium prausnitzii</i> - No change in <i>Oscillospira</i> Med diet compared to baseline: ↓ <i>Prevotella</i> ↑ <i>Roseburia</i> ↑ <i>Oscillospira</i> ↑ <i>Parabacteroides distasonis</i>	- After 1 y: no differences in main metabolic variables (glucose, HbA1c, insulin sensitivity index, TG, TC, HDL, LDL) between groups ↑ Insulin sensitivity index for both the LFHCC and Med diets, when measured from an OGTT performed at basal time and after 1 year of dietary intervention
Pu et al. 2014 [28]	N=25 (Finished per diet with stool sample: N=9-17) F: 76% 1 stool sample after interventions	Adults with at least one cardiovascular risk factor Mean (SD) Age: 53.6 (11.7) y BMI: 29.6 (4.59) kg/m ²	Randomized, controlled, double-blind, crossover clinical trial Country: Canada • 7-day rotation iso-caloric menu (3 meals, 2 snacks, 3000 kcal/d: CHO: 50E% Protein: 15E% Fat: 35E% • 60 g/d dietary oils equally distributed to 2 beverage shakes at breakfast and supper • Five oil treatments	Oil treatments (all diets were low in SFA): High MUFA, E%: IG1: canola oil [Canola; 63% MUFA, 20% LA, 10% ALA] IG2: DHA enriched canola-oil [CanolaDHA; 64% MUFA, 13% LA, 6% DHA] IG3: high OA canola oil (CanolaOleic; 72% MUFA, 15%	Comparisons between groups (no information on baseline microbiota): - Oil treatments had no significant impact on richness (Chao1, ACE) and α -diversity (Shannon, Simpson) - β -diversity did not change among treatments - Phylum distribution did not fluctuate across treatments or among MUFA vs PUFA groups - Average ratio of <i>Bacteroidetes</i> -to- <i>Firmicutes</i> was 0.15 across diets and did not differ among interventions - Genera <i>Parabacteroides</i> , <i>Prevotella</i> , <i>Turicibacter</i> , and family <i>Enterobacteriaceae</i> were positively correlated to MUFA-rich diets, while genus <i>Isobaculum</i> was correlated to PUFA-rich diets (R ² =0.43, Q ² =0.07) - CanolaDHA correlated to family <i>Lachnospiraceae</i>	- BMI had no significant impact on richness (Chao1, ACE) and α -diversity (Shannon, Simpson) - Rarefaction curves showed higher richness and diversity in OW/OB compared to NW participants - Similarity/ differences in microbiota among treatments and BMI (β -diversity) were compared using PCoA and PERMANOVA analyses of Bray-Curtis distances: Difference in OW vs OB ↑ Proportion of <i>Firmicutes</i> in OB compared to the combined NW/OW group - At the genus level, PLS-DA analysis confirmed a significant difference in the composition of bacteria among three BMI groups (R ² =0.60, Q ² =0.32) - TG was negatively correlated with phylum <i>Aquificae</i>

			Each treatment phase lasted 30 days, separated with 4 weeks washout periods	<p>LA, 2% ALA)</p> <p>High PUFA, E%: IG4 a blend of corn oil/safflower oil (CornSaff; 18% MUFA, 69% LA) – high n6 PUFA</p> <p>IG5: a blend of flax oil/safflower oil (FlaxSaff; 18% MUFA, 38% LA, 32% ALA) – high n3 PUFA</p>	<p>and phylum <i>Firmicutes</i> whereas CanolaOleic was associated with genera <i>Faecalibacterium</i> and <i>Coprobacillus</i> ($R^2=0.78$, $Q^2=0.45$)</p> <p>- CornSaff (but not FlaxSaff) had an impact on genera <i>Eggerthella</i>, <i>Slackia</i>, <i>Soehngenia</i>, <i>Anaerostipes</i>, <i>Robinsoniella</i>, <i>Phascolarctobacterium</i> ($R^2=0.67$, $Q^2=0.22$)</p> <p>In OW participants:</p> <p>- The genera <i>Streptococcus</i>, <i>Tepidimicrobium</i>, <i>Robinsoniella</i>, and <i>Turicibacter</i> were correlated to MUFA-rich and <i>Coriobacterium</i> and <i>Mogibacterium</i> to PUFA-rich diets ($R^2=0.69$, $Q^2=0.26$)</p> <p>- Comparing CanolaDHA and CanolaOleic, the genera <i>Adlercreutzia</i>, <i>Coriobacterium</i>, <i>Alistipes</i>, and <i>Robinsoniella</i> were correlated with CanolaDHA and <i>Lactobacillus</i> with CanolaOleic ($R^2=0.90$, $Q^2=0.60$)</p> <p>- Comparing PUFA-rich diets, CornSaff was associated with the genus <i>Adlercreutzia</i> and FlaxSaff with the genera <i>Collinsella</i>, <i>Barnesiella</i>, <i>Streptococcus</i>, <i>Roseburia</i>, <i>Coprobacillus</i>, and the family <i>Peptostreptococcaceae</i> ($R^2=0.98$, $Q^2=0.74$)</p> <p>In obese participants:</p> <p>- The genera <i>Parabacteroides</i>, <i>Prevotella</i>, <i>Flexithrix</i>, and <i>Fusibacter</i>; the family <i>Enterobacteriaceae</i>, and phylum <i>Firmicutes</i> were correlated to MUFA-rich diets, but no specific taxa was associated with PUFA-rich diets ($R^2=0.66$, $Q^2=-0.20$)</p> <p>- Comparing CanolaDHA and CanolaOleic, only the genus <i>Parasutterlla</i> correlated with CanolaDHA ($R^2=0.91$, $Q^2=0.29$)</p> <p>- Comparing the PUFA-rich diets, the genera <i>Collinsella</i>, <i>Hydrogenobaculum</i>, and <i>Parabacteroides</i> were impacted by the CornSaff, while the genus <i>Clostridium</i> was correlated to the FlaxSaff diet ($R^2=0.98$, $Q^2=0.63$)</p>	<p>($r=-0.27$) but positively with <i>Cyanobacteria</i> ($r=0.24$)</p> <p>- LDL was positively correlated with phylum <i>Proteobacteria</i> ($r=0.28$)</p> <p>- HDL was positively correlated with <i>Verrucomicrobia</i> ($r=0.21$)</p> <p>- In CanolaDHA treatment, TC levels positively correlated with <i>Firmicutes</i> ($r=0.55$)</p> <p>- In CornSaff treatment, TC levels were correlated with <i>Bacteroidetes</i> ($r=0.64$) and <i>Bacteroidetes</i>-to-<i>Firmicutes</i> ratio ($r=-0.65$)</p>
Rajkumar et al. 2014 [29]	N=60 F: 50%	OW, healthy adults aged 40-60 y Mean (range) Age: 49 (40-60) y BMI: 28.8 (27-30) kg/m ²	Randomized, placebo-controlled trial Country: India Fecal samples were obtained at baseline and after 45 days (6 weeks of intervention)	Participants received either (1) placebo (CG) (2) VSL#3 capsules (not considered here) (3) n3 PUFA capsules providing 180 mg EPA and 120 mg DHA per	n3 group: No effect on gut microbiota	At baseline: - Participants with vs without lipid abnormalities had lower total <i>lactobacilli</i> , <i>bifidobacteria</i> , and <i>streptococcus</i> and higher <i>Escherichia coli</i> and <i>bacteroides</i> - Similar trend for persons with vs without insulin resistance CG (compared to baseline):

				<p>day (IG) (4) n3 PUFA capsule + VSL#3 (not considered here)</p> <p>Intervention effects are only considered for n3 PUFA (IG) compared to CG</p>		<p>↑ FBG rose slightly</p> <p>n3 group (compared to baseline): ↓ insulin levels and FBG ↓ TC, TG, LDL, VLDL ↑ HDL, atherogenic index</p>
--	--	--	--	---	--	--

199 ↓ reduced ↑ increased

200

201

202

203

204

205

206

207 Table 2: Characteristics and results of the observational studies included in the systematic review

Study	Population		Study design and exposure		Results / Associations with outcomes	
Reference	N	Characteristics	Description	Fat intake	Microbiota	Metabolic health
Brahe et al. 2015 [30]	N=53 F: 100%	Postmenopausal obese women (BMI 30-45 kg/m ²) Mean (SD) Age: 60 (6) y BMI: 34.5 (3.8) kg/m ²	Cross-sectional study Country: Denmark Baseline assessment of a study sample recruited for a dietary intervention study 3-d weighed dietary intake Fecal sample	Dietary intake/d, mean (SD): Total energy, kj: 7572 (1797) E%: Fat: 35.3 (6.3) CHO: 40.8 (6.8) Protein: 18.9 (3.6) Fiber, g: 21.3 (6.0)	<i>Faecalibacterium prausnitzii</i> A2-165 and <i>Bacteroides pectinophilus</i> which were associated with a healthy metabolic profile were negatively correlated with E% of fat intake (r=-0.47 and r=-0.32, resp.). <i>Akkermansia muciniphila</i> which was associated with a healthy lipid profile was negatively associated with E% of fat intake (r=-0.28). <i>Clostridium bolteae</i> which was associated with an unhealthy metabolic profile was positively associated with E% of fat intake (r=0.35).	Negative correlation between metabolic markers of insulin resistance and the bacterial species <i>Bacteroides faecis</i> , <i>Intestinibacter bartlettii</i> , <i>Bifidobacterium longum</i> , <i>Faecalibacterium prausnitzii</i> A2-165, <i>Dorea longicatena</i> . The negative correlation between <i>Faecalibacterium prausnitzii</i> A2-165 and markers of insulin resistance disappeared after adjustment for fat intake. Positive correlation between metabolic markers of insulin resistance and the bacterial species <i>Ruminococcus torques</i> , <i>Clostridium bolteae</i> , <i>Eubacterium ramulus</i> , <i>Bilophila wadsworthia</i> Association between a healthy serum lipid profile and the following bacterial species: <i>Odoribacter splanchnicus</i> , <i>Bacteroides pectinophilus</i> , <i>Bacteroides cellulosilyticus</i> , <i>Bacteroides nordii</i> , <i>Roseburia inulinivorans</i> , <i>Akkermansia muciniphila</i> , <i>Faecalibacterium prausnitzii</i> A2-165, and <i>Bifidobacterium longum</i> Association between an unhealthy serum lipid profile and the bacterial species <i>Catenibacterium mitsuokai</i> , and <i>Holdemanella bififormis</i>
Fernandes et al. 2014 [31]	N=94 LN N=52 F: 57.7% OW/OB N=42 F: 50%	NW, OW and OB adults Mean (SEM) LN (BMI ≤25 kg/m ²): Age 32.0 (1.8) y BMI: 21.8 (0.3) kg/m ² Asian: 44% Caucasian: 50% Black: 2% Hispanic: 4% OW/OB (BMI >25)	Cross-sectional study Case-control study: Comparison of LN vs OW/OB group Country: Canada 3-d diet record Fecal sample	Dietary intake/d, mean (SEM), LN / OW/OB: Energy, kcal: 2035 (80) / 2063 (101) E%: Fat: 34 (1) / 36 (1) CHO: 47.3 (1.2) / 45.0 (1.4) Protein: 17 (1) / 18 (1) g/1000 kcal: SFA: 11.6 (0.5) / 12.6 (0.7) MUFA: 11.3 (0.6) 12.0 (0.7)	Combined groups: - Intake of PUFA was negatively correlated with <i>Bacteroidetes</i> (r=-0.21), all bacteria (r=-0.22) and <i>Firmicutes</i> (r=-0.25)	- F/B ratio was not different between the groups - LN (compared with OW/OB): ↑ <i>Escherichia coli</i> - BMI was inversely related to the number of <i>Bacteroidetes</i> (r=-0.21) and <i>Escherichia coli</i> (r=-0.34) - No association between the BMI and the log <i>Firmicutes</i> -to- <i>Bacteroides/Prevotella</i> ratio - No differences in the proportion of participants between groups who were <i>Archaea</i> positive

		kg/m ²): Age: 37.9 (2.0) BMI: 30.3 (0.7) kg/m ² Asian: 31% Caucasian: 55% Black: 12% Hispanic: 2%		PUFA, 5.2 (0.3) / 6.0 (0.5) Total fiber 11 (1) / 10 (1) Alcohol 1.4 (0.6) / 1.7 (0.5) TC: 129 (9) / 139 (9) Trans FA, g/d: 0.76 (0.12) / 0.76 (0.15)		
Mayorga Reyes et al. 2016 [37]	N=9 F: 66.7% N=3 in each group, LN, OW, OB	Young adults Mean (SD) Age: 27.1 (6.27) LN: BMI: 19.8 (0.94) kg/m ² WC: 67.7 (1.53) cm OW: BMI: 27.2 (0.51) kg/m ² WC: 87.8 (8.22) cm OB: BMI: 41.3 (5.25) kg/m ² WC: 114.3 (2.31) cm OW and OB persons had a slightly higher intake of SFA than recommended	Cross-sectional study Country: Mexico Semi-quantitative FFQ Phyla and bacterial species from fecal samples	Average dietary intake/d of LN / OW / OB: Energy, kcal: 2688 / 2520 / 1667 Fiber, g: 30.6 / 22.9 / 18.7 Range (E%): Fat: 36-40 CHO: 46.2-52.8 Protein: 11.2-14.8 SFA: 9-11 USFA: 8-9 No difference in the intake of SFA and USFA among the groups	- No correlation between food intake and abundance of <i>Firmicutes</i> and <i>Bacteroidetes</i> phyla or between food intake and <i>Bacteroides thetaiotaomicron</i> , <i>Faecalibacterium prausnitzii</i> , <i>Clostridium leptum</i> or <i>Prevotella</i> - Abundance of <i>Bifidobacterium longum</i> was positively correlated with an intake of foods that contained USFA - Intake of fiber was correlated to the abundance of the <i>Bacteroidetes</i> phylum	- No differences in the abundance of the phylum <i>Bacteroidetes</i> among groups - Abundance of <i>Firmicutes</i> in LN and OW groups was two logarithmic order of magnitude (LOM) greater than in OB <i>Prevotella</i> and <i>Bacteroides thetaiotaomicron</i> were not different among groups - LN and OW participants had one LOM more of <i>Faecalibacterium prausnitzii</i> than OB participants - LN and OW had four LOM greater <i>Clostridium leptum</i> abundance than did the OB group - Abundance of <i>Bifidobacterium longum</i> in the LN was two LOM more than in OW and five LOM more than in OB
Nakayama et al. 2017 [38]	N=43 Ormoc N=19 F: 36.8% Baybay N=24 F: 41.7%	7-9 year old children Mean (SD) Ormoc city (urban): Age: 8.11 (0.66) y BMI: 18.8 (4.5) kg/m ² Baybay city (rural): Age: 8.21 (0.51) y BMI: 14.8 (1.4) kg/m ²	Cross-sectional study Country: Philippines (Leyte island) FFQ for dietary assessment Fecal sample 85 and 95th percentiles were	Dietary intake/d, mean (SD) Ormoc city (urban): E%: Fat: 26.8 (5.2) CHO: 60.4 (6.0) Protein: 12.9 (2.3) g/d: SFA: 29.2 (10.7) MUFA: 22.0 (8.1) PUFA 8.94 (4.24) Trans FA, mg: 0.34 (0.22)	Ormoc city (compared to Baybay city): ↑ <i>Bacteroidaceae</i> ↑ <i>Ruminococcaceae</i> Baybay city (compared to Ormoc city): ↑ <i>Prevotellaceae</i> - No differences in <i>Bifidobacteriaceae</i> and <i>Lachnospiraceae</i> between cities Positive correlation of fat intake with: - <i>Firmicutes</i> -to- <i>Bacteroidetes</i> (F/B) ratio - <i>Firmicutes</i>	- All OW/OB children were living in Ormoc, suggesting a link between OW/OB and modern high-fat dietary habits - Higher fat intake in the OW/OB group than in the NW/UW group - F/B ratio was higher and relative abundance of <i>Prevotella</i> was lower in the OW/OB than in the NW/UW group (observed power determined retrospectively was not statistically high enough to warrant significance)

			used for the classification into OW and OB groups, resp. Participants below 15th percentile were classified as underweight.	Baybay city (rural): E%: Fat: 17.9 (4.7) CHO: 71.6 (6.0) Protein: 11.2 (2.2) g/d: SFA: 15.9 (5.9) , MUFA: 13.5 (5.7) PUFA: 4.69 (1.63) Trans FA, mg: 4.72 (18.84)	<ul style="list-style-type: none"> - an <i>Oscillibacter</i> sp. - a series of <i>Bacteroides/ Parabacteroides</i> spp. - genus <i>Bacteroides</i> - Order <i>Clostridiales</i> <p>Negative correlation of fat intake with:</p> <ul style="list-style-type: none"> - <i>Bacteroidetes</i> - family <i>Prevotellaceae</i> / genus <i>Prevotella</i> - genus <i>Succinivibrio</i> 	- The correlation between altered gut microbiota and high BMI suggests that a high-fat diet associated obesity is present among Filipino children on Leyte island
Org et al. 2017 [36]	N=531 F: 0%	45-70 year-old men Mean (SD) Age: 61.97 (5.45) y BMI: 27.92 (3.60) kg/m ²	Cross-sectional analysis Data based on a follow-up study of the population-based study cohort (subcohort of the METSIM cohort) Country: Finland Fecal samples	No information on dietary fat intake Serum metabolites of fatty acids	<ul style="list-style-type: none"> - Several associations with various fatty acids, accounting altogether for 41% of all taxonomy level (19 out of 46) and 33.8% of all OTU level (51 out of 151) associations <p>The most significant associations were observed with the abundance of members of the genus <i>Blautia</i> and phylum <i>Tenericutes</i>:</p> <ul style="list-style-type: none"> - Abundance of <i>Blautia</i> was positively associated with SFA and MUFA and negatively associated with degree of unsaturation and PUFA, including n3, DHA, n6, and LA - Negative associations of SFA with phylum <i>Tenericutes</i> - Negative associations of MUFA with <i>Peptococcaceae</i> - Positive associations of PUFA with <i>Tenericutes</i> and <i>Peptococcaceae</i> (for the latter also with LA, n6 PUFA) - Positive association of n3 PUFA incl. DHA with <i>Bacteroidales</i> 	<ul style="list-style-type: none"> - Fasting glucose levels were strongly associated with unclassified <i>Coriobacteriaceae</i> and several OTUs from <i>Blautia</i> were positively associated with pyruvate and glycerol - Higher abundances of genus <i>Methanobrevibacter</i> (<i>Archaea</i>), <i>Tenericutes</i>, <i>Peptococcaceae</i> and <i>Christensenellaceae</i> correlated with lower TG levels - No differences in either bacterial richness or in the F/B ratio between participants with different body weights and predisposition to T2D <p>In persons with high BMI:</p> <ul style="list-style-type: none"> ↑ the family <i>Tissierellaceae</i> and the genus <i>Blautia</i> ↓ <i>Archaea</i> (<i>Methanobrevibacter</i>) ↑ the genus <i>Anaerostipes</i> <p>In pre-diabetic persons:</p> <ul style="list-style-type: none"> ↓ lower abundances of an OTU from the families <i>Ruminococcaceae</i> and <i>Christensenellaceae</i> and the genus <i>Methanobrevibacter</i> - ↑ abundance of the order <i>Bacteroidales</i> in obese subjects was associated with lower HOMA and the higher abundance of the genus <i>Collinsella</i> with higher levels of glycerol and phenylalanine - Opposite effect in LN subjects
Röytiö et al. 2017 [32]	N=100 (88 with complete data) F: 100%	OW/OB women at early pregnancy (≤17 week of gestation) Mean (SD) Age: 30.1 (4.7) y	Cross-sectional analysis within an ongoing mother–infant dietary intervention trial Country: Finland	Group 1: low-fiber/moderate-fat group (N=57) - fiber intake (<25 g/d) / total fat intake (25-40 E%) Group 2: high-	<ul style="list-style-type: none"> - Intakes of total fat and different fat types (except for n3 PUFA) were negatively associated with one (PUFA, n6 PUFA) or more indicators of gut microbiota diversity and richness (α-diversity, measured as Chao1, observed OTU, phylogenetic diversity, Shannon index) - SFA were negatively associated with all diversity 	<ul style="list-style-type: none"> - Contradictory findings were found at the genus level within the family <i>Lachnospiraceae</i>: <i>Lachnospira</i> was negatively and <i>Blautia</i> positively correlated with concentrations of various sized VLDL particles and TG in VLDL - The genus <i>Lachnospira</i> was negatively associated with serum TG

		<p>Pre-pregnancy BMI: 30.2 (4.6) kg/m²</p> <p>The three groups did not differ in BMI</p>	<p>A 10-h fasting blood sample was drawn from the participants.</p> <p>Fecal samples from mothers were collected.</p> <p>3 d food diaries recorded within the week before the study visit</p>	<p>fiber/moderate fat group (N=18) - fiber (≥ 25 g/d) / total fat intake (25-40 E%) but higher energy intake than the other groups.</p> <p>Group 3: low-fiber/high-fat group (N=13) - fat intake (≥ 40E%). SFA consumption above reference (>10 E%) Consumption of SFA, MUFA and PUFA, in addition to total fat, was higher than in the other groups; consumption of fiber and total CHO lower than recommended.</p>	<p>and richness indexes, whereas n3 PUFA showed no correlation</p> <p>- Negative correlations between the intake of fat (E%) and SFA (E%) and relative abundance in the family <i>Barnesiellaceae</i></p> <p>Comparison of the 3 groups, Mean (SD): Higher in the high-fiber/moderate-fat group compared with the low-fiber/high-fat group: α-diversity (Chao1 index) 406.2 (44.4) vs 341.0 (SD 57.9), phylogenetic diversity (PD) 39.0 (4.5) vs 31.3 (6.7) and observed number of OTU 355.8 (38.7) vs 293.8 (59.0). The low fiber/moderate-fat group did not differ from the other groups (Chao 1 index 380.0 (57.3), PD 35.8 (5.9), observed number of OTU 333.0 (55.2).</p>	<p>- The genus <i>Blautia</i> was positively associated with VLDL diameter, but negatively with the diameters of LDL and HDL.</p> <p>- No correlations were detected between gut microbiota richness indexes and serum lipidomics variables.</p> <p>- No differences were detected in markers of low-grade inflammation, serum lipidomic variables or zonulin concentration among the three diet groups.</p>
<p>Simoes et al. 2013 [33]</p>	<p>N=40 F: 55%</p> <p>NW: N=11</p> <p>OW: N=18</p> <p>OB: N=11</p>	<p>Monozygotic twin pairs</p> <p>Mean (SD) NW: Age: 26 (3) y BMI: 22.9 (2.2) kg/m²</p> <p>OW: Age: 29 (3) y BMI: 26.5 (1.2) kg/m²</p> <p>OB: Age: 28 (4) y BMI: 32.4 (2.1) kg/m²</p>	<p>Cross-sectional analysis of data from a population-based longitudinal survey</p> <p>Country: Finland</p> <p>Participants were divided into 3 BMI groups: NW (19\leqBMI<25) OW (25\leqBMI<30) OB (BMI\geq30)</p> <p>3-d food diary, supervised by a specialist</p>	<p>NW / OW / OB, mean (SD): Energy, MJ: 8.0 (1.7) / 8.4 (2.2) / 9.8 (2.0)</p> <p>g/d: Fat: 77 (29) / 75 (26) / 85 (22) SFA: 30 (12) / 28 (9.9) / 32 (8.6) MUFA: 23 (8.9) / 19 (6.9) / 23 (6.9) PUFA: 10 (4.0) / 10 (5.1) / 13 (5.2) n3 PUFA: 1.8 (0.7) / 1.5 (0.7) / 1.6 (0.6) n6 PUFA: 7.9 (3.2) / 8.6 (4.5) / 11 (4.4)</p> <p>CHO: 200 (50) / 219 (58) / 255 (51) Protein: 85 (29) / 86 (31) / 81 (31) Total fiber: 21 (14) / 16 (6.1) / 17 (5.5)</p>	<p>High energy intake compared to lower intake: ↓ <i>Bacteroides</i> spp. ↑ <i>Bifidobacteria</i></p> <p>Greater MUFA compared to lower consumption: ↓ <i>Bifidobacteria</i></p> <p>- Increased ingestion of n3 PUFA had a significant association with higher numbers of bacteria within the <i>Lactobacillus</i> group</p> <p>- Greater n6 PUFA consumption was negatively correlated with the numbers of <i>bifidobacteria</i></p> <p>- Co-twins with the same SFA intake had very similar <i>Bacteroides</i> spp. profiles (80-100% similarity) whereas the twin pairs with distinct SFA intake had low similarity (0-25%). The group of co-twins who consumed similar amounts of fiber had very low <i>bifidobacterial</i> similarity (0-25%).</p>	<p>- The numbers of bacteria within the different bacterial groups, as measured by qPCR, did not differ between BMI groups.</p> <p>- The diversity of the studied bacterial groups, defined as the number of the bands obtained by different group-specific PCR DGGE did not differ between BMI groups.</p> <p>- No relation was found between the intrapair DGGE profile similarities and the co-twin concordance for BMI, intrapair difference in BMI, or body fat.</p>

Yamaguchi et al. 2016 [35]	N=59 F: 57.6% NW: N=42 OW: N=13 OB: N=4	T2D patients Median (IQ) Age: 65 (58.5-69.0) y BMI: 23.0 (20.4-25.6) kg/m ²	Cross-sectional study Country: Japan Fasting blood and fecal samples Data-based short FFQ	Dietary intake/d, mean (SD): Energy: 1692 (380) kcal E%: Fat: 23.2 (5.3) CHO: 57.5 (5.2) Protein: 13.2 (2.2)	- <i>Clostridium</i> cluster IV: negatively correlated with fat intake (r=-0.261); positively correlated with CHO (r=0.266) - <i>Clostridium</i> cluster XI: positively correlated with both fat (r=0.301) and protein intake (r=0.363)	- Bifidobacterium spp., order Lactobacillales and <i>Bacteroides</i> spp. were negatively correlated with fasting blood glucose (r=-0.264) - <i>Clostridium</i> subcluster XIVa: positively correlated with TC (r=0.385) - <i>Clostridium</i> cluster IV was negatively correlated with fecal acetate which was shown to be beneficial for glucose tolerance - Propionate and acetate were negatively correlated to insulin and HOMA. Butyrate was positively correlated to HDL. Total SCFA were negatively correlated with insulin and HOMA.
Yang et al. 2017 [34]	N=71 F: 100% Low: N=24 Moderate: N=23 High: N=24	Premenopausal women aged 19-49 y Mean (95% CI) Low fitness: Age: 40.4 (36.9-44.0) y BMI: 31.7 (30.2-33.1) kg/m ² BF%: 40.6 38.1-43.0) Moderate fitness: Age: 39.7 (35.5-43.8) y BMI: 27.9 (26.7-29.1) kg/m ² BF%: 35.5 (33.2-37.8) High fitness: Age: 30.6 (25.6-35.6) y BMI: 24.6 (23.0-26.2) kg/m ² BF%: 28.0 (25.0-31.0)	Cross-sectional study Country: Finland Food diary records 3 groups according to cardiorespiratory fitness (tertiles of VO2max): (1) high fitness (high) (2) moderate fitness = control (moderate) (3) low fitness (low)	Daily mean intake of low/ moderate/ high fitness group (E%, unless otherwise stated): Fat: 32.8/ 34.1/ 35.2 CHO: 47.1/ 45.6/ 44.7 Protein: 18.4/ 18.0/ 18.0 Alcohol (E%): 1.73/ 2.38/ 0.63 Fiber, g/d: 20.4/ 24.1/ 21.0	High fitness (low BMI) compared to low fitness group (high BMI): ↑ proportions of <i>Bacteroides</i> ↓ <i>EreC</i> , phylum <i>Firmicutes</i> No differences between groups for <i>Bifidobacterium</i> , <i>Enterobacteria</i> , <i>Faecalibacterium prausnitzii</i> - <i>EreC</i> was positively correlated with fat intake (r=0.258) - <i>EreC</i> was inversely correlated with CHO intake (r=-0.252)	- <i>EreC</i> was positively correlated with BF% (r=0.382) and TG (r=0.390) and negatively with HDL (r=0.26) After adjustment for BF%, correlations disappeared. - Multivariable regression analysis showed that <i>EreC</i> contributed the most to VO2max, BF%, Leptin, HDL, TG - VO2max was negatively correlated with BF% (r=0.755), TG (r=-0.274) and leptin (r=-0.574)

↓ lower ↑ higher

208

209

210 Risk of bias assessment*211 Quality assessment of randomized controlled interventional studies based on the Cochrane tool*

212 In total, six RCT were evaluated based on the Cochrane risk of bias tool (Table 3). All six studies were
213 considered of 'low-risk' with regard to 'random sequence generation'. The sequence generation was
214 described as being computer generated in each of the studies, with the exception of Rajkumar et al.
215 2014 [29] where an identification number was assigned to each participant by a scientist blind to the
216 treatments corresponding with each code. Four of the six studies were considered 'low-risk' with
217 regard to 'allocation concealment' as the assignment of codes was reported as blinded. In two
218 studies [26, 27], risk was considered 'unclear' as the allocation of codes was not reported. The
219 blinding of participants was considered 'low-risk' in two of the six studies as blinding of participants
220 and study coordinators was performed for each study [25, 28]. Four of the remaining studies were
221 considered 'high-risk' as either the participants or study group were not reported to be blinded. The
222 blinding of outcomes was considered 'low-risk' in five of the six studies, with the exception of one
223 study [26] which was considered 'unclear' as the analysis performed in the study was not clearly
224 reported as blinded. With regard to 'incomplete outcomes', two of the six studies were considered
225 'low-risk' as the number of missing data points was minimal and would not be considered as a source
226 of bias in these studies. One study [28] was considered 'high-risk' as a number of participants did not
227 provide a sample at every time point throughout the study, and three of the remaining studies were
228 considered 'unclear' as missing data points and/or the final number of participants was not stated.
229 Overall, all six studies were considered 'low-risk' with regard to 'selective reporting' as all outcome
230 assessments provided in the methods were stated in the results section, and two of the six studies
231 were considered 'high-risk' with regard to other forms of bias due to gender (all participants were
232 men) [27] and the obese state of the participants [28].

233

234 Quality assessment of case-control studies based on Newcastle-Ottawa Scale

235 In total, nine cross-sectional studies were analyzed according to the NOS (Table 4). Two studies
236 received a score of 2 in the 'selection' category because of their study design. Three studies received

237 a score of 3 in the 'selection' category as they successfully completed the criteria required for an
238 adequate case-study definition with the selection of suitable control groups. Four studies received
239 the maximum of 4 stars in the 'selection' category as they completed all of the necessary
240 requirements for the selection and definition of a high quality case-control study. In the
241 'comparability' category six studies received a score of 2 (maximum score) due to the number of
242 variable confounding factors which were included and adjusted for in the analysis, such as BMI, age
243 and dietary intake among others. Four of the studies received a score of 1 in this 'comparability'
244 category as only one confounding factor was controlled for throughout the study. In the 'exposure'
245 category six studies received a score of 2 (maximum score is 4) as the methods used to attain the
246 results for each study did not appear to create potential bias in either the case or control groups;
247 however these six studies did not clearly describe the non-response rate in each group and therefore
248 were not awarded an additional star. Two studies received a score of 1 in the 'exposure' category as
249 the ascertainment of results was only adequate in either the case or control group. Fernandes et al.
250 2014 received a total score of 3 in the 'exposure' category as only one participant was reported to
251 drop-out of the study. Overall, three studies received a total score of 8, three studies received a total
252 score of 7, one study received a total score of 6 and two studies received a total score of 5 in the NOS
253 quality assessment scale. Studies which received higher scores of 7 or 8 indicate better
254 methodological quality.

255

256

257

258

259 Table 3: Risk of bias assessment of the randomized controlled trials (RCT) based on Cochrane risk of
260 bias tool. Risk of bias of each item was judged as low (+), high (-) or unclear (?).

Reference	Random sequence generation	Allocation concealment	Blinding of participants /personnel	Blinding of outcome assessment	Incomplete outcome	Selective reporting	Other bias
Balfego et al. 2016 [24]	+	+	-	+	?	+	+
Blaedel et al. 2016 [25]	+	+	+	+	+	+	+
Fava et al. 2013 [26]	+	?	-	?	?	+	+
Haro et al. 2016 [27]	+	?	-	+	+	+	-
Pu et al. 2014 [28]	+	+	+	+	-	+	-
Rajkumar et al. 2014 [29]	+	+	-	+	?	+	+

262 Table 4: Quality assessment for the selected studies based on Newcastle-Ottawa Quality Assessment Scale for case-control studies. Total score ranges from 0 to
 263 9. Higher scores indicated better methodological quality.

Reference	SELECTION				COMPARABILITY		EXPOSURE		TOTAL
	Case definition	Representativeness of the cases	Selection of controls	Definition of controls	Study controls for...	Ascertainment of the exposure	Same method of ascertainment for cases and controls	Non-response rate	
Brahe et al. 2015 [30]	*	*	*		**	*	*		7
Fernandes et al. 2014 [31]	*	*	*		*	*	*	*	7
Mayorga Reyes et al. 2016 [37]	*	*	*		*	*	*		6
Nakayama et al. 2017 [38]	*	*	*	*	*	*	*		7
Simoes et al. 2013 [33]	*	*	*	*	**	*	*		8
Org et al. 2017 [36]	*	*	*	*	**	*	*		8
Röytiö et al. 2017 [#] [32]		*	*		**		*		5
Yamaguchi et al. 2016 [#] [35]	*	*			**	*			5
Yang et al. 2017 [34]	*	*	*	*	**	*	*		8

264 [#] Analysis of associations between diet and the gut microbiota composition and clinical markers only, no case-control design

265

266

267

268

269 **High versus low fat diets in relation to the intestinal microbiota and metabolic outcomes**

270 *Interventional studies*

271 High fat interventional diets reduced total bacteria compared to baseline, whereas this reduction
272 was not seen in low fat/high carbohydrate diets in adults at increased risk of MetS. Instead, the latter
273 increased *Bifidobacterium spp.* and *Bacteroides spp.* At the end of the interventions with three high-
274 fat and two low-fat diets, no differences in cardiometabolic risk factors were observed [26].
275 Accordingly, no differences between either a high-fat Mediterranean (high in MUFA) or a low-fat diet
276 were observed in main metabolic variables of glucose/insulin metabolism and lipoprotein profile
277 after one year of intervention. The Mediterranean diet resulted in a decrease of the genus *Prevotella*
278 and increased the genera *Roseburia* and *Oscillospira* and the species *Parabacteroides distasonis*
279 compared to baseline [27].

280

281 *Observational studies*

282 In overweight and obese pregnant women with different dietary patterns the intake of total fat was
283 negatively associated with gut microbiota diversity and richness [32]. In a cross-sectional study with
284 postmenopausal obese women, a healthy lipoprotein profile showed a positive association with the
285 species *Faecalibacterium prausnitzii* A2-165, *Bacteroides pectinophilus* and *Akkermansia muciniphila*
286 which were negatively correlated with fat intake. *Clostridium bolteae* was positively correlated with
287 fat intake and showed a positive correlation with markers of insulin resistance [30]. In Philippine
288 children, the families *Bacteroidaceae* and *Ruminococcaceae* were higher in urban living children with
289 low fat intake compared to rural living children with very low fat intake who had higher abundance of
290 the family *Prevotellaceae*. Fat intake was positively correlated with the *Firmicutes*-to-*Bacteroidetes*
291 (*F/B*) ratio, *Firmicutes*, an *Oscillibacter* species, various *Bacteroides/ Parabacteroides* species, genus
292 *Bacteroides* and the order *Clostridiales* (*Firmicutes*) and was negatively correlated with the genera
293 *Bacteroidetes* and *Prevotella* (family *Prevotellaceae*) and *Succinivibrio* (phylum *Proteobacteria*) [38].
294 Contrary correlations with fat intake were seen for *Clostridium* cluster XI which correlated positively
295 and *Clostridium* cluster IV (*Clostridium leptum*) which correlated negatively in 59 patients with type 2

296 diabetes. The latter was also negatively correlated with fecal acetate which was shown to be
297 beneficial for glucose tolerance [35]. In premenopausal women, *Eubacterium rectale-Clostridium*
298 *coccoides (EreC)* was positively correlated with fat intake and showed a positive correlation with
299 body fat percentage. In a multivariable regression analysis *EreC* contributed the most to body fat
300 percentage, HDL and TG [34].

301

302 **High versus low SFA diets in relation to the intestinal microbiota and metabolic outcomes**

303 *Interventional studies*

304 In a crossover RCT in healthy men who received for 21 days either a diet enriched with whole-fat milk
305 (40 E% of fat) which contains mainly SFA or an isoenergetic standard diet with 35 E% of fat no effects
306 on the fecal microbiota, on the blood lipoprotein profile or on insulin and glucose concentrations
307 were observed [25]. A high total fat/high SFA diet (18 E% of SFA, 38 E% of total fat) increased
308 *Faecalibacterium prausnitzii* compared to baseline. The comparison of five diets including the high
309 SFA diet, two isoenergetic low SFA diets (10 E%) with the same amount of total fat and two low
310 fat/high carbohydrate diets did not result in differences in BMI, waist circumference, body fat
311 percentage, blood pressure and insulin sensitivity parameters between the diets at the end of the
312 intervention [26].

313

314 *Observational studies*

315 In overweight and obese pregnant women, SFA consumption was negatively associated with all gut
316 microbiota diversity and richness indexes [32]. In adult men, higher abundance of the genus *Blautia*
317 which was positively associated with SFA serum metabolites was detected in persons with high BMI.
318 Higher abundance of the phylum *Tenericutes* which was negatively associated with SFA metabolites
319 correlated with lower triglyceride levels [36]. In a study with monozygotic twin pairs, co-twins with
320 the same SFA intake had very similar *Bacteroides spp.* profiles whereas low similarity was observed in
321 twin pairs with distinct SFA intake [33].

322

323 **High MUFA diets in relation to the intestinal microbiota and metabolic outcomes**

324 *Interventional studies*

325 Two high fat/high MUFA diets decreased total bacterial numbers compared to a low fat/high
326 carbohydrate diet and compared to baseline. While waist circumference decreased in the high MUFA
327 group with low glycemic index compared to baseline, no significant changes in BMI, waist
328 circumference, body fat percentage, blood pressure or insulin sensitivity between the different
329 MUFA- and/or PUFA-rich diets were detected at the end of the intervention [26]. In a cross-over RCT
330 with identical total fat intake but different MUFA-rich oil treatments, high MUFA diets showed no
331 effect on richness/diversity indexes, the phylum distribution or *Bacteroidetes*-to-*Firmicutes* ratio.
332 MUFA-rich diets were positively correlated to the genera *Parabacteroides*, *Prevotella* and
333 *Turicibacter*, and the family *Enterobacteriaceae*. The BMI had no significant association with richness
334 (Chao1, ACE) and α -diversity (Shannon, Simpson) although rarefaction curves showed higher richness
335 and diversity in overweight/obese compared to normal weight participants. Additionally, similarity
336 and differences in microbiota among BMI (β -diversity) showed differences in normal weight versus
337 obese participants. Also a higher proportion of the phylum *Firmicutes* was reported in obese
338 compared to the combined normal weight/overweight group. Triglyceride levels were negatively
339 correlated with the phylum *Aquificae* and positively with *Cyanobacteria* while LDL was positively
340 correlated with *Proteobacteria* and HDL with *Verrucomicrobia* [28]. Compared to baseline, a MUFA-
341 rich Mediterranean diet decreased the genus *Prevotella* and increased the genera *Roseburia* and
342 *Oscillospira*, and the species *Parabacteroides distasonis* while a low-fat diet with a high proportion of
343 complex carbohydrates (LFHCC) showed opposite effects in the genera *Prevotella* and *Roseburia*, no
344 effect on *Oscillospira*, and an increase of the species *Faecalibacterium prausnitzii*. While insulin
345 sensitivity was increased after one year on both diets compared to baseline, no differences in the
346 main metabolic outcomes of glucose/insulin status and lipoprotein profile were observed between
347 the groups [27].

348

349 *Observational studies*

350 In adult men, abundance of the genus *Blautia* which was shown to be positively associated with
351 MUFA serum metabolites was increased in persons with high BMI. Higher abundance of the phylum
352 *Tenericutes* which was negatively associated with MUFA metabolites correlated with lower
353 triglyceride levels. MUFA were also negatively associated with the family *Peptococcaceae* [36]. In a
354 study with monozygotic twin pairs, higher compared to lower MUFA consumption was correlated to
355 lower number of the genus *Bifidobacterium*. The numbers of bacteria within the different bacterial
356 groups as measured by qPCR and diversity of studied bacterial groups did not differ between BMI
357 groups [33].

358

359 **High PUFA diets in relation to the intestinal microbiota and metabolic outcomes**

360 *Interventional studies*

361 Three RCT investigated the effects of n3 PUFA enriched diets on the gut microbiota and found no
362 effects on the intestinal microbiota compared to control groups [24, 28, 29]. However, only one of
363 the interventions had a longer duration of six months [24], whereas the other two interventions
364 lasted for only 30 days [28] or six weeks [29]. Ingestion of a docosahexaenoic acid (DHA)-enriched
365 high MUFA diet and a high n3 (α -linolenic acid, ALA) or n6 (linoleic, LA) PUFA diet had no impact on
366 bacterial richness, diversity or phylum distribution. Compared to a high MUFA diet with low n3 PUFA
367 a DHA-enriched high MUFA diet correlated to the family *Lachnospiraceae* and the phylum *Firmicutes*.
368 Total cholesterol levels were positively associated with *Firmicutes* in the group with the DHA-
369 enriched high MUFA diet. In the n6 LA enriched diet total cholesterol levels were positively
370 correlated with the phylum *Bacteroidetes* and negatively with the *Bacteroidetes*-to-*Firmicutes* ratio
371 [28].

372

373 *Observational studies*

374 In accordance with the results of the interventional studies, R yti  et al. (2017) also reported no
375 correlation between any diversity or richness index and n3 PUFA intake in pregnant women [32]. In
376 adults, an inverse association between PUFA intake and *Bacteroidetes*, all bacteria and *Firmicutes*

377 was shown. The BMI was inversely related to the number of *Bacteroidetes* [31]. Org et al. (2017) [36]
378 investigated serum metabolites of fatty acids in 45-70 year-old men and found that the abundance of
379 the genus *Blautia* was negatively associated with PUFA including n6 and n3 PUFA and was increased
380 in participants with higher BMI. In contrast positive associations with PUFA were observed with the
381 genus *Bacteroidales*, the phylum *Tenericutes* and the family *Peptococcaceae*. Higher abundances of
382 the latter two correlated with lower triglyceride levels [36]. In monozygotic twin pairs high n3 PUFA
383 ingestion was associated with higher numbers of bacteria within the *Lactobacillus* group whereas
384 higher intake of n6 PUFA was negatively associated with the abundance of the genus *Bifidobacterium*
385 [33]. In contrast, in a study with nine participants the abundance of the species *Bifidobacterium*
386 *longum* was positively correlated with the intake of unsaturated fatty acids and was higher in lean
387 than in overweight and obese participants [37].

388

389 **Discussion**


390 To our knowledge, this is the first systematic review that compiles and provides effects/associations
391 of dietary fat quantity and quality on/with the gut microbiota composition and cardiometabolic
392 health in humans. Based on 15 included studies, our results of interventional trials do not suggest
393 strong effects of dietary fat quantity or quality on the gut microbiota or on metabolic health
394 outcomes while observational studies indicate associations with the gut microbiota and health
395 outcomes. It has to be noted that half of the interventional studies had a relatively short duration of
396 three to six weeks [25, 28, 29], which may be one reason why they showed no strong effects of fat
397 type on either the gut microbiota or on metabolic health. Figure 2 gives an overview of the main
398 results of intervention and observational studies included in this systematic review. As evidence
399 provided by observational studies is less strong than that from intervention studies the following
400 discussion section will primarily focus on the latter studies.

401 It should be noted that the value of the results of microbiota analysis is limited due to the use of
402 qPCR and FISH methods in most of the studies published, which do not allow a complete taxonomic
403 assessment of the hundreds of species inhabiting the intestine. Consequently, comprehensive

404 analyses by Next Generation Sequencing methods are required to better reflect the impact of fat
 405 quantity and quality on gut microbiota at the community structure level. Additionally, a recent study
 406 conducted in three different populations that investigated the temporal stability of specific
 407 microbiome features, based on 16S ribosomal RNA (rRNA) gene profiles and including two biological
 408 samples from each subject separated by approximately six months, revealed a large variability and
 409 low temporal stability of major phyla and alpha-diversity metrics. This makes it very difficult to draw
 410 reliable conclusions from cross-sectional studies as well as to identify robust associations between
 411 the microbiota changes with health outcomes in intervention studies unless several samples are
 412 analyzed longitudinally [39].

413

414



415

416 Figure 2: Main associations between dietary fat and intestinal microbiota and between intestinal
 417 microbiota and metabolic health markers. Both, interventional (A) and observational studies (B)
 418 show associations between total fat intake, mainly SFA, and reduction of bacterial abundance,

419 diversity and richness in the gut. (A) Dietary fat interventions do not suggest strong effects on gut
420 microbiota. (B) High intake of total fat or SFA is positively correlated with the abundance of
421 *Clostridium bolteae* and *Blautia* respectively, both species associated with unhealthy metabolic
422 outcomes (insulin resistance and increased BMI). PUFA-enriched diet is associated with increased
423 abundance of *Tenericutes* which is associated with lower levels of TG in plasma.

424 ^a In the study by Org et al. 2017 [36], serum levels instead of dietary intake were measured.

425

426

427 *High fat, high SFA and high MUFA diets*

428 High fat (Western) diets have been shown to be associated with lower richness and diversity of the
429 intestinal microbiota in animals and humans [40-42] whereas high intake of vegetables and fruit (rich
430 in fiber) is associated with high richness and diversity [43, 44]. This is in line with the observed
431 reduction of total bacteria after dietary interventions with high compared to low fat content in adults
432 with increased MetS risk [26]. Also in pregnant women total fat and SFA intake were negatively
433 associated with the gut microbiota richness and diversity [32]. While previous studies showed that
434 lower microbiome richness is associated with obesity, higher fat mass, insulin resistance and
435 dyslipidemia compared to higher richness [45, 46], no significant differences of metabolic markers
436 after high versus low fat interventional diets were observed in a randomized study although total
437 bacteria decreased after the high fat interventions [26]. Accordingly, in adults with at least one
438 cardiovascular risk factor the BMI had no significant impact on richness and α -diversity although
439 rarefaction curves showed higher microbiota richness and diversity in overweight/obese compared
440 to normal weight participants. Additionally, similarity and differences in microbiota among BMI (β -
441 diversity) showed differences in normal weight versus obese participants [28].

442 In contrast to SFA, results on the effects of MUFA-rich diets are less consistent. In mice, MUFA do not
443 seem to affect microbiota richness and diversity [47] and may even increase bacterial density [48].

444 Also in adults with increased cardiovascular disease risk, microbiome richness and diversity were not

445 affected by high MUFA intake after an interventional period of 30 days [28]. However, in adults with
446 an increased risk of MetS, high MUFA diets decreased total bacteria cells after 24 weeks of
447 intervention compared to a high carbohydrate diet and compared to baseline. As this decrease was
448 not accompanied by a decrease in any of the fluorescence in-situ hybridization (FISH)-enumerated
449 bacteria, unrecognized bacterial populations must have been reduced which may suggest that the
450 high MUFA diets negatively affected richness and diversity of the gut microbiota [26]. Also in a cross-
451 sectional study, a high MUFA intake was negatively associated with microbiota diversity and richness
452 [32].

453 A higher relative abundance of *Firmicutes* and a lower abundance of *Bacteroidetes* after the ingestion
454 of high-fat diets was previously reported in mice [41, 49, 50] and was confirmed in a study with
455 children indicating that fat intake is positively correlated with *Firmicutes* and the *F/B* ratio but
456 negatively with *Bacteroidetes* [38]. A higher proportion of *Firmicutes* was also reported in obese
457 compared to the combined normal weight/overweight group in adults with at least one
458 cardiovascular disease risk factor [28]. Results of the included intervention and observational studies
459 in general confirm previous findings suggesting that a decrease of *Bacteroidetes* and an increase of
460 *Firmicutes* are correlated with obesity in humans [51] and animals [6, 47, 52]. Nevertheless, a recent
461 meta-analysis pooling data of 10 studies conducted by 16S rRNA gene sequencing did not confirm
462 such association [53]. Animal studies indicate that changes in the gut microbiota composition are
463 directly caused by fat intake rather than the degree of obesity because contrary to a high fat/SFA
464 diet, a high fat/MUFA diet was not associated with changes in the gut microbiota but resulted in a
465 higher degree of obesity than an energy-matched low-fat/SFA diet. However, differences in the gut
466 microbiota composition were only found on the high fat/SFA diet and, thus, seem to result from the
467 overflow of dietary fat but not from the obese phenotype [47]. Another study reported consistent
468 and strong changes in the gut microbiota composition upon switching to a high fat diet for both wild-
469 type and RELM β (expression depends upon the presence of the gut microbiome) Knockout (KO) mice

470 indicating that the high fat diet itself but not the obese state caused the alterations of the microbiota
471 [50].

472 In contrary to the reported results with high fat diets, high MUFA diets ingested for 30 days showed
473 no effect on the *Bacteroidetes*-to-*Firmicutes* ratio in an RCT with different oil treatments but identical
474 energy% of total fat [28] which confirms previous results in mice [47].

475 Intervention studies also showed changes in single bacterial genus or species by high or low fat diets
476 although there was no consistent trend. Supplementary Table 2 provides a short summary
477 description of the affected bacteria.

478 High abundance of the genus *Prevotella* is typical for a high carbohydrate and fiber rich diet [8, 43,
479 54]. Accordingly, a high-fat Mediterranean diet resulted in a decrease of the genus *Prevotella*
480 compared to a low-fat/high complex carbohydrate diet and compared to baseline in obese coronary
481 heart disease patients but did not result in differences in metabolic endpoints [27]. In an RCT with
482 identical total fat intake of 35 energy%, MUFA-rich diets were correlated with the genus *Prevotella*
483 and the composition of bacteria differed between different weight status groups [28]. This was also
484 the case in a study with pregnant women which reported a higher relative abundance of the genus
485 *Prevotella* and of the family *Prevotellaceae* in obese than in overweight women [55].

486 Higher abundance of the species *Faecalibacterium prausnitzii* was observed in association with high-
487 fiber diets and with beneficial effects on intestinal barrier function [56], on the fat-free mass as seen
488 in young male children [57] and on health [58]. In contrast, according to results of included studies, a
489 diet high in total fat (and SFA) negatively affects *Faecalibacterium prausnitzii* [26, 30] whereas a low-
490 fat/high complex carbohydrate diet increased *Faecalibacterium prausnitzii* compared to a high fat
491 Mediterranean diet but both improved insulin sensitivity [27] which may have resulted from higher
492 vegetable and fiber intake typical for these diets. Despite an increase of *Faecalibacterium prausnitzii*
493 in a high fat/high SFA group, an RCT did not detect changes in adiposity or cardiometabolic risk

494 factors [26] whereas an observational study reported positive associations of *Faecalibacterium*
495 *prausnitzii* with a healthy lipoprotein profile [30].

496 Pu et al. showed a positive correlation between MUFA-rich diets and populations of
497 *Enterobacteriaceae* which is the only family in the order *Enterobacteriales* [28]. In contrast, in mice
498 fed a high-fat diet supplemented with a MUFA-rich (oleic acid) compound decreased the order
499 *Enterobacteriales* and *Clostridium* cluster XIVa which had been increased by the high-fat diet and
500 increased *Bifidobacterium* spp. which had been decreased by the high-fat diet [48].

501

502 *High PUFA diets*

503 In contrast to high fat and high SFA diets, evidence from three included RCT suggests that n3 PUFA-
504 enriched diets have no effect on the gut microbiota compared to control diets [24, 28, 29] although
505 two of the interventions lasted for only 30 [28] and 45 [29] days, respectively. The above null effects
506 of dietary PUFA were also evidenced in the cross-over intervention study performed in the frame of
507 the MyNewGut project where no impact on gut microbiota, anthropometry, metabolism, and
508 physiology was observed after administration of fish oil capsules containing 3.6 g/d n3 PUFA (DHA
509 and eicosapentanoic acid, EPA) [59]. In contrast, a recently published cross-over intervention with 4
510 g/d n3 PUFA for 8 weeks did not find changes in α or β diversity, or phyla composition but showed
511 an increased abundance of beneficial bacteria such as *Bifidobacterium*, *Roseburia* and *Lactobacillus*
512 [60]. Thus, a high dose and long-duration intake of n3 PUFA may be necessary to induce positive
513 effects on the microbiome composition. The beneficial effects on metabolic outcomes observed by
514 n3 PUFA-rich diets in the two included studies above [24, 29] which confirm previous studies with n3
515 PUFA supplementation [61, 62] seem to be independent of the gut microbiota.

516 Also in mice microbial diversity was not affected by diets high in PUFA [47]. Accordingly, diets
517 including n3 PUFA-enriched (DHA) high MUFA oil and high n3 (ALA) or n6 (LA) PUFA oil treatments
518 for 30 days did not result in changes in bacterial richness, diversity or phylum distribution [28].

519 A DHA-enriched high MUFA oil treatment for 30 days correlated to *Firmicutes* which were positively
520 associated with total cholesterol levels [28]. This type of diet also correlated to *Lachnospiraceae* [28]
521 which has been reported to be increased by high fat diets in animal studies [41, 63]. Compared to
522 diets rich in SFA, diets rich in n3 or n6 PUFA resulted in lower decreases in *Bacteroidetes* in mice [64].

523

524 *Limitations*

525 The *reduction* of one of the major components of diet usually influences the ingestion of other
526 macronutrients. In this regard, an increase of dietary fat intake is mostly paralleled by lower
527 carbohydrate and fiber consumption. Therefore, it is hardly possible to attribute observed changes
528 only to fat or specific fatty acids if there is no comparison group with identical intake of the other
529 nutrients. This increases the risk of bias in observational studies and in the RCT comparing a
530 Mediterranean and a low-fat diet [27]. Also, the energy intake can vary because of different fat
531 intake and can influence the results. Some papers indicated that the energy content of the diet is as
532 important as or even more important than the composition of the diet in driving gut microbiota
533 changes [30-33]. Thus, associations between dietary fat/fatty acid intake and the intestinal
534 microbiota as well as between the microbiota and metabolic health outcomes reported from
535 observational studies may have been influenced by other dietary factors and energy intake as well.
536 Also, other lifestyle factors (e.g. physical activity) influence an individuals' microbiota and (metabolic)
537 health and may have affected the results of the included studies [65]. Another limitation, particularly
538 of observational studies is that – with the exception of the study on biomarkers [36] – fat and fatty
539 acid intake was estimated based on participant's self-reported dietary intake. Further, most studies
540 included only small sample sizes of highly selected participants.

541

542 **Conclusions and recommendations**

543 Based on the included intervention and observational studies, this systematic review indicates that a
544 high fat diet and a high fat diet rich in SFA may exert unfavorable effects on the gut microbiota
545 characterized by lower richness and diversity and is generally associated with an unhealthy metabolic

546 state. Results on diets rich in MUFA are less consistent. MUFA may have no effect on gut microbiota
547 richness and diversity or may negatively affect total bacterial numbers and gut microbiota richness
548 and diversity. In contrast, diets rich in n3 or n6 PUFA do not seem to negatively affect the gut
549 microbiota or metabolic health outcomes. Thus, high fat intake and in particular high SFA intake
550 should be reduced in favor of higher PUFA intake. Considering the conflicting results and a potential
551 negative effect of MUFA on the gut microbiome, the dietary recommendation to reduce SFA and to
552 replace them with (plant-sources of) MUFA and PUFA [66, 67] may need additional research.
553 However, data are not consistent and the overall evidence was weak due to risk of bias and small,
554 not representative samples. Additional ongoing data analyses within the MyNewGut project will help
555 to elucidate the role of the diet in altering the gut microbiota and associations with metabolic health
556 outcomes. In particular, high quality longitudinal and intervention studies comparing effects of SFA,
557 MUFA and specific n3 and n6 PUFA are missing.

558

559 **Acknowledgement**

560 We would like to thank Lara Christianson for her valuable support in the systematic literature search
561 and related presentation and Rieke Baumkötter for her support in further literature research and
562 administration.

563

564 **Funding**

565 The MyNewGut project is financially supported by a grant from the EU 7th Framework Programme
566 under Grant Agreement 613979. The EU is not liable for the content presented in this publication.

567

568

569 Supplementary Table 1: Medline (PubMed) search strategy

570

Search	Terms
#1	"Gastrointestinal Microbiomes*" [Title/Abstract] OR "Gut Microflora*" [Title/Abstract] OR "Gastrointestinal Flora*" [Title/Abstract] OR "Gut Flora*" [Title/Abstract] OR "Gastrointestinal Microbiota*" [Title/Abstract] OR "Gastrointestinal Microflora*" [Title/Abstract] OR "Enteric Bacteria*" [Title/Abstract] OR "Intestinal Microbiome*" [Title/Abstract] OR "Intestinal Microbiota*" [Title/Abstract] OR "Intestinal Microflora*" [Title/Abstract] OR "gut microbiome*" [Title/Abstract] OR "gut microbiota*" [Title/Abstract] OR "gut bacteria*" [Title/Abstract]
#2	"Gastrointestinal Microbiome" [Mesh]
#3	#1 OR #2
#4	"dietary fat*" [Title/Abstract] OR "fatty acid*" [Title/Abstract] OR "unsaturated fatty acid*" [Title/Abstract] OR "saturated fatty acid*" [Title/Abstract] OR "polyunsaturated fatty acid*" [Title/Abstract] OR "omega-3 fatty acid*" [Title/Abstract] OR "omega-6 fatty acid*" [Title/Abstract] OR "eicosapentaenoic acid*" [Title/Abstract] OR "docosahexaenoic acid*" [Title/Abstract] OR "arachidonic acid*" [Title/Abstract] OR "linoleic acid*" [Title/Abstract] OR "linolenic acid*" [Title/Abstract] OR "oleic acid*" [Title/Abstract] OR "stearic acid*" [Title/Abstract] OR "palmitic acid*" [Title/Abstract] OR "fish oil*" [Title/Abstract] OR "fat intake*" [Title/Abstract]
#5	"Fatty Acids" [MeSH Terms] OR "Dietary Fats" [MeSH Terms]
#6	#4 OR #5
#7	"metabolic syndrome*" [Title/Abstract] OR "metabolic health*" [Title/Abstract] OR "abdominal obesity*" [Title/Abstract] OR "blood pressure*" [Title/Abstract] OR "blood sugar*" [Title/Abstract] OR "serum triglyceride*" [Title/Abstract] OR "hypertension*" [Title/Abstract] OR "hyperglycemia*" [Title/Abstract] OR "dyslipidemia*" [Title/Abstract] OR "insulin resistance*" [Title/Abstract] OR "insulin resistant*" [Title/Abstract] OR "obesity*" [Title/Abstract] OR "overweight*" [Title/Abstract] OR "adiposity*" [Title/Abstract] OR "adipositas*" [Title/Abstract] OR "HOMA IR*" [Title/Abstract] OR "cardiovascular disease*" [Title/Abstract] OR "cardiovascular syndrome*" [Title/Abstract] OR "type 2 diabetes*" [Title/Abstract] OR "cholesterol*" [Title/Abstract] OR "LDL cholesterol*" [Title/Abstract] OR "HDL cholesterol*" [Title/Abstract] OR "triglycerides*" [Title/Abstract] OR "hypertriglyceridemia*" [Title/Abstract] OR "hypercholesterolemia*" [Title/Abstract] OR "hyperinsulinemia*" [Title/Abstract] OR "insulin*" [Title/Abstract] OR "serum glucose*" [Title/Abstract] OR "blood glucose*" [Title/Abstract] OR "HbA1c*" [Title/Abstract] OR "glycated hemoglobin*" [Title/Abstract] OR "waist circumference*" [Title/Abstract] OR "coronary artery disease*" [Title/Abstract] OR "stroke*" [Title/Abstract]
#8	Metabolic Syndrome [MeSH Terms] OR Obesity [MeSH Terms] OR Cardiovascular Diseases [MeSH Terms] OR Diabetes Mellitus, Type 2 [MeSH Terms] OR Cholesterol [MeSH Terms] OR Stroke [MeSH Terms]
#9	#7 OR #8
#10	#3 AND #6 AND #9
#11	#3 AND #6 AND #9 AND Filters: Humans

571

572

573 Supplementary Table 2: Description of bacterial genus/species influenced by high/low fat diets in the
574 included intervention studies

575

Change by fat intake (Energy%)	Bacteria genus or species	Description [44, 68]
<ul style="list-style-type: none"> - Increase after high fat (38 E%, SFA: 18 E%) [26] - Increase after low fat (28 E%, SFA: 8 E%)/ high carbohydrate [27] 	<i>Faecalibacterium prausnitzii</i>	Phylum: <i>Firmicutes</i> Class: <i>Clostridia</i> Order: <i>Clostridiales</i> Family: <i>Clostridiaceae</i> Genus: <i>Faecalibacterium</i> <ul style="list-style-type: none"> - Species of grampositive bacteria - Associated with plant-based diets high in carbohydrate - Associated with anti-inflammatory properties - Buyrate producer
<ul style="list-style-type: none"> - Increase after low fat (28 E%)/ high carbohydrate / decrease after high fat Med (35 E%) [27] 	<i>Prevotella</i>	Phylum and Class: <i>Bacteroidetes</i> Order: <i>Bacteroidales</i> Family: <i>Prevotellaceae</i> <ul style="list-style-type: none"> - Genus of gramnegative bacteria - Associated with plant-based high carbohydrate and high fiber diets - Propionate producer
<ul style="list-style-type: none"> - Increase after low fat (28 E%)/ high carbohydrate / decrease after high fat Med (35 E%) [27] 	<i>Roseburia</i>	Phylum: <i>Firmicutes</i> Class: <i>Clostridia</i> Order: <i>Clostridiales</i> Family: <i>Lachnospiraceae</i> <ul style="list-style-type: none"> - Genus of grampositive bacteria - Associated with plant-based diets high in carbohydrates - Buyrate producer
<ul style="list-style-type: none"> - Increase after low fat (28 E%)/ high carbohydrate [26] 	<i>Bifidobacterium</i> spp.	Phylum and class: <i>Actinobacteria</i> Order: <i>Bifidobacteriales</i> Family: <i>Bifidobacteriaceae</i> Genus: <i>Bifidobacterium</i> <ul style="list-style-type: none"> - Species of grampositive bacteria - Associated with fiber, particularly fructooligosaccharide intake
<ul style="list-style-type: none"> - Increase after low fat (28 E%)/ high carbohydrate [26] 	<i>Bacteroides</i> spp.	Phylum: <i>Bacteroidetes</i> Class: <i>Bacteroidia</i> Order: <i>Bacteroidales</i> Family: <i>Bacteroidaceae</i> Genus: <i>Bacteroides</i> <ul style="list-style-type: none"> - Species of gramnegative bacteria - Able to adapt to fiber rich and animal-based diets rich in protein and fat

576

578 **References**

- 579 1. Arora T, Backhed F. The gut microbiota and metabolic disease: current understanding and
580 future perspectives. *Journal of internal medicine*. 2016. Epub 2016/04/14. doi: 10.1111/joim.12508.
581 PubMed PMID: 27071815.
- 582 2. Cummings JH, Macfarlane GT. Role of intestinal bacteria in nutrient metabolism. *JPEN Journal*
583 *of parenteral and enteral nutrition*. 1997;21(6):357-65. Epub 1997/12/24. doi:
584 10.1177/0148607197021006357. PubMed PMID: 9406136.
- 585 3. Pihl AF, Fonvig CE, Stjernholm T, Hansen T, Pedersen O, Holm JC. The Role of the Gut
586 Microbiota in Childhood Obesity. *Childhood obesity (Print)*. 2016. Epub 2016/04/09. doi:
587 10.1089/chi.2015.0220. PubMed PMID: 27058515.
- 588 4. Pereira SS, Alvarez-Leite JI. Low-Grade Inflammation, Obesity, and Diabetes. *Current obesity*
589 *reports*. 2014;3(4):422-31. Epub 2015/12/03. doi: 10.1007/s13679-014-0124-9. PubMed PMID:
590 26626919.
- 591 5. Korpela K, de Vos WM. Antibiotic use in childhood alters the gut microbiota and predisposes
592 to overweight. *Microbial cell (Graz, Austria)*. 2016;3(7):296-8. Epub 2017/03/31. doi:
593 10.15698/mic2016.07.514. PubMed PMID: 28357367; PubMed Central PMCID: PMC5354595.
- 594 6. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated
595 gut microbiome with increased capacity for energy harvest. *Nature*. 2006;444(7122):1027-31. Epub
596 2006/12/22. doi: 10.1038/nature05414. PubMed PMID: 17183312.
- 597 7. Rampelli S, Candela M, Turroni S, Biagi E, Pflueger M, Wolters M, et al. Microbiota and
598 lifestyle interactions through the lifespan. *Trends in Food Science & Technology*. 2016. doi:
599 <http://dx.doi.org/10.1016/j.tifs.2016.03.003>.
- 600 8. Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, et al. Linking long-term
601 dietary patterns with gut microbial enterotypes. *Science (New York, NY)*. 2011;334(6052):105-8.
602 Epub 2011/09/03. doi: 10.1126/science.1208344. PubMed PMID: 21885731; PubMed Central PMCID:
603 PMC3368382.
- 604 9. Alcock J, Lin HC. Fatty acids from diet and microbiota regulate energy metabolism.
605 *F1000Research*. 2015;4(F1000 Faculty Rev):738. Epub 2015/01/01. doi:
606 10.12688/f1000research.6078.1. PubMed PMID: 27006755; PubMed Central PMCID:
607 PMC4797936.
- 608 10. Shen W, Gaskins HR, McIntosh MK. Influence of dietary fat on intestinal microbes,
609 inflammation, barrier function and metabolic outcomes. *The Journal of nutritional biochemistry*.
610 2014;25(3):270-80. Epub 2013/12/21. doi: 10.1016/j.jnutbio.2013.09.009. PubMed PMID: 24355793.
- 611 11. Kaliannan K, Wang B, Li XY, Bhan AK, Kang JX. Omega-3 fatty acids prevent early-life antibiotic
612 exposure-induced gut microbiota dysbiosis and later-life obesity. *International journal of obesity*
613 (2005). 2016;40(6):1039-42. Epub 2016/02/16. doi: 10.1038/ijo.2016.27. PubMed PMID: 26876435.
- 614 12. Tonucci LB, Olbrich Dos Santos KM, Licursi de Oliveira L, Rocha Ribeiro SM, Duarte Martino
615 HS. Clinical application of probiotics in type 2 diabetes mellitus: A randomized, double-blind, placebo-
616 controlled study. *Clinical nutrition (Edinburgh, Scotland)*. 2017;36(1):85-92. Epub 2016/01/07. doi:
617 10.1016/j.clnu.2015.11.011. PubMed PMID: 26732026.
- 618 13. Valentini L, Pinto A, Bourdel-Marchasson I, Ostan R, Brigidi P, Turroni S, et al. Impact of
619 personalized diet and probiotic supplementation on inflammation, nutritional parameters and
620 intestinal microbiota - The "RISTOMED project": Randomized controlled trial in healthy older people.
621 *Clinical nutrition (Edinburgh, Scotland)*. 2015;34(4):593-602. Epub 2014/12/03. doi:
622 10.1016/j.clnu.2014.09.023. PubMed PMID: 25453395.
- 623 14. Karlsson FH, Tremaroli V, Nookaew I, Bergstrom G, Behre CJ, Fagerberg B, et al. Gut
624 metagenome in European women with normal, impaired and diabetic glucose control. *Nature*.
625 2013;498(7452):99-103. Epub 2013/05/31. doi: 10.1038/nature12198. PubMed PMID: 23719380.
- 626 15. Yassour M, Lim MY, Yun HS, Tickle TL, Sung J, Song YM, et al. Sub-clinical detection of gut
627 microbial biomarkers of obesity and type 2 diabetes. *Genome medicine*. 2016;8(1):17. Epub

- 628 2016/02/18. doi: 10.1186/s13073-016-0271-6. PubMed PMID: 26884067; PubMed Central PMCID:
629 PMCPMC4756455.
- 630 16. Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, et al. A metagenome-wide association study of gut
631 microbiota in type 2 diabetes. *Nature*. 2012;490(7418):55-60. Epub 2012/10/02. doi:
632 10.1038/nature11450. PubMed PMID: 23023125.
- 633 17. Lin HV, Frassetto A, Kowalik EJ, Jr., Nawrocki AR, Lu MM, Kosinski JR, et al. Butyrate and
634 propionate protect against diet-induced obesity and regulate gut hormones via free fatty acid
635 receptor 3-independent mechanisms. *PloS one*. 2012;7(4):e35240. Epub 2012/04/17. doi:
636 10.1371/journal.pone.0035240. PubMed PMID: 22506074; PubMed Central PMCID:
637 PMCPMC3323649.
- 638 18. Zhang X, Shen D, Fang Z, Jie Z, Qiu X, Zhang C, et al. Human gut microbiota changes reveal the
639 progression of glucose intolerance. *PloS one*. 2013;8(8):e71108. Epub 2013/09/10. doi:
640 10.1371/journal.pone.0071108. PubMed PMID: 24013136; PubMed Central PMCID:
641 PMCPMC3754967.
- 642 19. Schwiertz A, Taras D, Schafer K, Beijer S, Bos NA, Donus C, et al. Microbiota and SCFA in lean
643 and overweight healthy subjects. *Obesity (Silver Spring, Md)*. 2010;18(1):190-5. Epub 2009/06/06.
644 doi: 10.1038/oby.2009.167. PubMed PMID: 19498350.
- 645 20. den Besten G, Havinga R, Bleeker A, Rao S, Gerding A, van Eunen K, et al. The short-chain
646 fatty acid uptake fluxes by mice on a guar gum supplemented diet associate with amelioration of
647 major biomarkers of the metabolic syndrome. *PloS one*. 2014;9(9):e107392. Epub 2014/09/10. doi:
648 10.1371/journal.pone.0107392. PubMed PMID: 25203112; PubMed Central PMCID:
649 PMCPMC4159349.
- 650 21. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews
651 and meta-analyses: the PRISMA statement. *PLoS medicine*. 2009;6(7):e1000097. Epub 2009/07/22.
652 doi: 10.1371/journal.pmed.1000097. PubMed PMID: 19621072; PubMed Central PMCID:
653 PMCPMC2707599.
- 654 22. Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions (The
655 Cochrane Collaboration)*. Chichester: John Wiley & Sons Ltd; 2011.
- 656 23. Wells GS, B., O'Connell DP, J., Welch VL, M., Tugwell P. *The Newcastle-Ottawa Scale (NOS) for
657 assessing the quality of nonrandomised studies in meta-analyses*. 2011.
- 658 24. Balfego M, Canivell S, Hanzu FA, Sala-Vila A, Martinez-Medina M, Murillo S, et al. Effects of
659 sardine-enriched diet on metabolic control, inflammation and gut microbiota in drug-naive patients
660 with type 2 diabetes: a pilot randomized trial. *Lipids Health Dis*. 2016;15:78. Epub 2016/04/20. doi:
661 10.1186/s12944-016-0245-0. PubMed PMID: 27090218; PubMed Central PMCID: PMCPMC4836051.
- 662 25. Blädel T, Holm JB, Sundekilde UK, Schmedes MS, Hess AL, Lorenzen JK, et al. A randomised,
663 controlled, crossover study of the effect of diet on angiotensin-like protein 4 (ANGPTL4) through
664 modification of the gut microbiome. *Journal of Nutritional Science*. 2016;5. doi: 10.1017/jns.2016.38.
- 665 26. Fava F, Gitau R, Griffin BA, Gibson GR, Tuohy KM, Lovegrove JA. The type and quantity of
666 dietary fat and carbohydrate alter faecal microbiome and short-chain fatty acid excretion in a
667 metabolic syndrome 'at-risk' population. *International journal of obesity (2005)*. 2013;37(2):216-23.
668 Epub 2012/03/14. doi: 10.1038/ijo.2012.33. PubMed PMID: 22410962.
- 669 27. Haro C, Montes-Borrego M, Rangel-Zuniga OA, Alcalá-Díaz JF, Gomez-Delgado F, Perez-
670 Martínez P, et al. Two Healthy Diets Modulate Gut Microbial Community Improving Insulin Sensitivity
671 in a Human Obese Population. *J Clin Endocrinol Metab*. 2016;101(1):233-42. Epub 2015/10/28. doi:
672 10.1210/jc.2015-3351. PubMed PMID: 26505825.
- 673 28. Pu S, Khazanehei HR, Krause DO, West SG, Kris-Etherton PM, Jenkins DJ, et al. Effects of
674 unsaturated fatty acids (USFA) on human gut microbiome profile in a subset of canola oil multicenter
675 intervention trial (COMIT). *FASEB Journal*. 2013;27.
- 676 29. Rajkumar H, Mahmood N, Kumar M, Varikuti SR, Challa HR, Myakala SP. Effect of probiotic
677 (VSL#3) and omega-3 on lipid profile, insulin sensitivity, inflammatory markers, and gut colonization
678 in overweight adults: a randomized, controlled trial. *Mediators Inflamm*. 2014;2014:348959. Epub
679 2014/05/06. doi: 10.1155/2014/348959. PubMed PMID: 24795503; PubMed Central PMCID:
680 PMCPMC3984795.

- 681 30. Brahe LK, Le Chatelier E, Prifti E, Pons N, Kennedy S, Hansen T, et al. Specific gut microbiota
682 features and metabolic markers in postmenopausal women with obesity. *Nutrition and Diabetes*.
683 2015;5. doi: 10.1038/nutd.2015.9.
- 684 31. Fernandes J, Su W, Rahat-Rozenbloom S, Wolever TMS, Comelli EM. Adiposity, gut
685 microbiota and faecal short chain fatty acids are linked in adult humans. *Nutrition and Diabetes*.
686 2014;4(JUNE). doi: 10.1038/nutd.2014.23.
- 687 32. Roytio H, Morkkala K, Vahlberg T, Laitinen K. Dietary intake of fat and fibre according to
688 reference values relates to higher gut microbiota richness in overweight pregnant women. *The*
689 *British journal of nutrition*. 2017;118(5):343-52. Epub 2017/09/14. doi:
690 10.1017/s0007114517002100. PubMed PMID: 28901891.
- 691 33. Simoes CD, Maukonen J, Kaprio J, Rissanen A, Pietilainen KH, Saarela M. Habitual dietary
692 intake is associated with stool microbiota composition in monozygotic twins. *J Nutr*.
693 2013;143(4):417-23. Epub 2013/01/25. doi: 10.3945/jn.112.166322. PubMed PMID: 23343669.
- 694 34. Yang Y, Shi Y, Wiklund P, Tan X, Wu N, Zhang X, et al. The association between
695 cardiorespiratory fitness and gut microbiota composition in premenopausal women. *Nutrients*.
696 2017;9(8). doi: 10.3390/nu9080792.
- 697 35. Yamaguchi Y, Adachi K, Sugiyama T, Shimozaoto A, Ebi M, Ogasawara N, et al. Association of
698 intestinal microbiota with metabolic markers and dietary habits in patients with type 2 diabetes.
699 *Journal of Gastroenterology and Hepatology (Australia)*. 2016;31:154. doi: 10.1111/jgh.13540.
- 700 36. Org E, Blum Y, Kasela S, Mehrabian M, Kuusisto J, Kangas AJ, et al. Relationships between gut
701 microbiota, plasma metabolites, and metabolic syndrome traits in the METSIM cohort. *Genome Biol*.
702 2017;18(1):70. Epub 2017/04/15. doi: 10.1186/s13059-017-1194-2. PubMed PMID: 28407784;
703 PubMed Central PMCID: PMC5390365.
- 704 37. Mayorga Reyes L, Gonzalez Vazquez R, Cruz Arroyo SM, Melendez Avalos A, Reyes Castillo
705 PA, Chavaro Perez DA, et al. Correlation between diet and gut bacteria in a population of young
706 adults. *Int J Food Sci Nutr*. 2016;67(4):470-8. Epub 2016/03/29. doi:
707 10.3109/09637486.2016.1162770. PubMed PMID: 27018166.
- 708 38. Nakayama J, Yamamoto A, Palermo-Conde LA, Higashi K, Sonomoto K, Tan J, et al. Impact of
709 westernized diet on gut microbiota in children on Leyte island. *Frontiers in Microbiology*.
710 2017;8(FEB). doi: 10.3389/fmicb.2017.00197.
- 711 39. Sinha R, Goedert JJ, Vogtmann E, Hua X, Porras C, Hayes R, et al. Quantification of Human
712 Microbiome Stability Over 6 Months: Implications for Epidemiologic Studies. *American journal of*
713 *epidemiology*. 2018;187(6):1282-90. Epub 2018/04/03. doi: 10.1093/aje/kwy064. PubMed PMID:
714 29608646; PubMed Central PMCID: PMC5982812.
- 715 40. Agus A, Denizot J, Thevenot J, Martinez-Medina M, Massier S, Sauvanet P, et al. Western diet
716 induces a shift in microbiota composition enhancing susceptibility to Adherent-Invasive *E. coli*
717 infection and intestinal inflammation. *Scientific reports*. 2016;6:19032. Epub 2016/01/09. doi:
718 10.1038/srep19032. PubMed PMID: 26742586; PubMed Central PMCID: PMC4705701.
- 719 41. Zhang C, Zhang M, Pang X, Zhao Y, Wang L, Zhao L. Structural resilience of the gut microbiota
720 in adult mice under high-fat dietary perturbations. *The ISME journal*. 2012;6(10):1848-57. Epub
721 2012/04/13. doi: 10.1038/ismej.2012.27. PubMed PMID: 22495068; PubMed Central PMCID:
722 PMC3446802.
- 723 42. Devkota S, Wang Y, Musch MW, Leone V, Fehlner-Peach H, Nadimpalli A, et al. Dietary-fat-
724 induced taurocholic acid promotes pathobiont expansion and colitis in *Il10*^{-/-} mice. *Nature*.
725 2012;487(7405):104-8. Epub 2012/06/23. doi: 10.1038/nature11225. PubMed PMID: 22722865;
726 PubMed Central PMCID: PMC3393783.
- 727 43. De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, et al. Impact of diet
728 in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa.
729 *Proceedings of the National Academy of Sciences of the United States of America*.
730 2010;107(33):14691-6. Epub 2010/08/04. doi: 10.1073/pnas.1005963107. PubMed PMID: 20679230;
731 PubMed Central PMCID: PMC2930426.

- 732 44. Benítez-Páez A, Gómez Del Pulgar EM, Kjolbaek L, Kirchner Brahe L, Astrup A, Larsen L, et al.
733 Impact of dietary fiber and fat on gut microbiota re-modeling and metabolic health. *Trends in Food*
734 *Science & Technology*. 2016;57:201-12.
- 735 45. Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, et al. Richness of human gut
736 microbiome correlates with metabolic markers. *Nature*. 2013;500(7464):541-6. Epub 2013/08/30.
737 doi: 10.1038/nature12506. PubMed PMID: 23985870.
- 738 46. Albenberg LG, Wu GD. Diet and the intestinal microbiome: associations, functions, and
739 implications for health and disease. *Gastroenterology*. 2014;146(6):1564-72. Epub 2014/02/08. doi:
740 10.1053/j.gastro.2014.01.058. PubMed PMID: 24503132; PubMed Central PMCID: PMC4216184.
- 741 47. de Wit N, Derrien M, Bosch-Vermeulen H, Oosterink E, Keshtkar S, Duval C, et al. Saturated
742 fat stimulates obesity and hepatic steatosis and affects gut microbiota composition by an enhanced
743 overflow of dietary fat to the distal intestine. *American journal of physiology Gastrointestinal and*
744 *liver physiology*. 2012;303(5):G589-99. Epub 2012/06/16. doi: 10.1152/ajpgi.00488.2011. PubMed
745 PMID: 22700822.
- 746 48. Mujico JR, Baccan GC, Gheorghe A, Diaz LE, Marcos A. Changes in gut microbiota due to
747 supplemented fatty acids in diet-induced obese mice. *The British journal of nutrition*.
748 2013;110(4):711-20. Epub 2013/01/11. doi: 10.1017/s0007114512005612. PubMed PMID:
749 23302605.
- 750 49. Turnbaugh PJ, Backhed F, Fulton L, Gordon JI. Diet-induced obesity is linked to marked but
751 reversible alterations in the mouse distal gut microbiome. *Cell host & microbe*. 2008;3(4):213-23.
752 Epub 2008/04/15. doi: 10.1016/j.chom.2008.02.015. PubMed PMID: 18407065; PubMed Central
753 PMCID: PMC3687783.
- 754 50. Hildebrandt MA, Hoffmann C, Sherrill-Mix SA, Keilbaugh SA, Hamady M, Chen YY, et al. High-
755 fat diet determines the composition of the murine gut microbiome independently of obesity.
756 *Gastroenterology*. 2009;137(5):1716-24.e1-2. Epub 2009/08/27. doi: 10.1053/j.gastro.2009.08.042.
757 PubMed PMID: 19706296; PubMed Central PMCID: PMC3687783.
- 758 51. Turnbaugh PJ, Hamady M, Yatsunencko T, Cantarel BL, Duncan A, Ley RE, et al. A core gut
759 microbiome in obese and lean twins. *Nature*. 2009;457(7228):480-4. Epub 2008/12/02. doi:
760 10.1038/nature07540. PubMed PMID: 19043404; PubMed Central PMCID: PMC2677729.
- 761 52. Ley RE, Backhed F, Turnbaugh P, Lozupone CA, Knight RD, Gordon JI. Obesity alters gut
762 microbial ecology. *Proceedings of the National Academy of Sciences of the United States of America*.
763 2005;102(31):11070-5. Epub 2005/07/22. doi: 10.1073/pnas.0504978102. PubMed PMID: 16033867;
764 PubMed Central PMCID: PMC1176910.
- 765 53. Sze MA, Schloss PD. Looking for a Signal in the Noise: Revisiting Obesity and the Microbiome.
766 *MBio*. 2016;7(4).
- 767 54. Yatsunencko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, et al. Human
768 gut microbiome viewed across age and geography. *Nature*. 2012;486(7402):222-7. Epub 2012/06/16.
769 doi: 10.1038/nature11053. PubMed PMID: 22699611; PubMed Central PMCID: PMC3376388.
- 770 55. Houttu N, Mokka K, Laitinen K. Overweight and obesity status in pregnant women are
771 related to intestinal microbiota and serum metabolic and inflammatory profiles. *Clinical nutrition*
772 (Edinburgh, Scotland). 2017. Epub 2018/01/18. doi: 10.1016/j.clnu.2017.12.013. PubMed PMID:
773 29338886.
- 774 56. Martin R, Miquel S, Chain F, Natividad JM, Jury J, Lu J, et al. *Faecalibacterium prausnitzii*
775 prevents physiological damages in a chronic low-grade inflammation murine model. *BMC*
776 *microbiology*. 2015;15:67. Epub 2015/04/19. doi: 10.1186/s12866-015-0400-1. PubMed PMID:
777 25888448; PubMed Central PMCID: PMC4391109.
- 778 57. Smith-Brown P, Morrison M, Krause L, Davies PSW. Male-specific Association Between Fat-
779 Free Mass Index and Fecal Microbiota in 2- to 3-Year-Old Australian Children. *Journal of Pediatric*
780 *Gastroenterology and Nutrition*. 2018;66(1):147-51.
- 781 58. Miquel S, Martin R, Rossi O, Bermudez-Humaran LG, Chatel JM, Sokol H, et al.
782 *Faecalibacterium prausnitzii* and human intestinal health. *Current opinion in microbiology*.
783 2013;16(3):255-61. Epub 2013/07/09. doi: 10.1016/j.mib.2013.06.003. PubMed PMID: 23831042.

- 784 59. Kjølbæk L, Benítez-Páez A, Gómez del Pulgar EM, Brahe LK, Liebisch G, Matysik S, et al.
785 Arabinoxylan oligosaccharides and polyunsaturated fatty acid effects on gut microbiota and
786 metabolic markers in overweight individuals with signs of metabolic syndrome: a randomized cross-
787 over trial. *Clinical nutrition (Edinburgh, Scotland)*. 2018;Under Review.
- 788 60. Watson H, Mitra S, Croden FC, Taylor M, Wood HM, Perry SL, et al. A randomised trial of the
789 effect of omega-3 polyunsaturated fatty acid supplements on the human intestinal microbiota. *Gut*.
790 2018;67(11):1974-83. Epub 2017/09/28. doi: 10.1136/gutjnl-2017-314968. PubMed PMID:
791 28951525.
- 792 61. Rimm EB, Appel LJ, Chiuve SE, Djousse L, Engler MB, Kris-Etherton PM, et al. Seafood Long-
793 Chain n-3 Polyunsaturated Fatty Acids and Cardiovascular Disease: A Science Advisory From the
794 American Heart Association. *Circulation*. 2018. Epub 2018/05/19. doi:
795 10.1161/cir.0000000000000574. PubMed PMID: 29773586.
- 796 62. Innes JK, Calder PC. The Differential Effects of Eicosapentaenoic Acid and Docosahexaenoic
797 Acid on Cardiometabolic Risk Factors: A Systematic Review. *International journal of molecular*
798 *sciences*. 2018;19(2). Epub 2018/02/10. doi: 10.3390/ijms19020532. PubMed PMID: 29425187;
799 PubMed Central PMCID: PMC5855754.
- 800 63. Lecomte V, Kaakoush NO, Maloney CA, Raipuria M, Huinao KD, Mitchell HM, et al. Changes in
801 gut microbiota in rats fed a high fat diet correlate with obesity-associated metabolic parameters.
802 *PloS one*. 2015;10(5):e0126931. Epub 2015/05/21. doi: 10.1371/journal.pone.0126931. PubMed
803 PMID: 25992554; PubMed Central PMCID: PMC4436290.
- 804 64. Liu T, Hougen H, Vollmer AC, Hiebert SM. Gut bacteria profiles of *Mus musculus* at the
805 phylum and family levels are influenced by saturation of dietary fatty acids. *Anaerobe*.
806 2012;18(3):331-7. Epub 2012/03/06. doi: 10.1016/j.anaerobe.2012.02.004. PubMed PMID:
807 22387300.
- 808 65. Rampelli S, Guenther K, Turroni S, Wolters M, Veidebaum T, Yiannis K, et al. Gut microbiome
809 is associated with diet, physical activity and health in children with or without excessive weight gain.
810 *Communications Biology*. 2018; in print.
- 811 66. EFSA Panel on Dietetic Products NaAN. Scientific Opinion on Dietary Reference Values for
812 fats, including saturated fatty acids, polyunsaturated fatty acids, monounsaturated fatty acids,
813 trans fatty acids, and cholesterol. *EFSA Journal*. 2010;8(3):1461 [107 pp.]. doi:
814 10.2903/j.efsa.2010.1461.
- 815 67. U.S. Department of Health and Human Services and U.S. Department of Agriculture (USDA).
816 2015-2020 Dietary Guidelines for Americans. 8th ed, December 2015.
- 817 68. Portune KJ, Benitez-Paez A, Del Pulgar EM, Cerrudo V, Sanz Y. Gut microbiota, diet, and
818 obesity-related disorders-The good, the bad, and the future challenges. *Molecular nutrition & food*
819 *research*. 2017;61(1). Epub 2016/06/12. doi: 10.1002/mnfr.201600252. PubMed PMID: 27287778.

820