ORIGINAL RESEARCH

Food Science & Nutrition

WILEY

Preservation of gallic acid and methyl gallate on purified Kilka fish oil oxidation by Rancimat

Maryam Asnaashari¹ | Reza Farhoosh² | Reza Farahmandfar¹

Revised: 15 May 2019

¹Department of Food Science and Technology, Sari Agricultural Sciences and Natural Resources University (SANRU), Sari, Iran

²Department of Food Science and Technology, Ferdowsi University of Mashhad (FUM), Mashhad, Iran

Correspondence

Reza Farahmandfar and Maryam Asnaashari, Department of Food Science and Technology, Sari Agricultural Sciences and Natural Resources University (SANRU), Sari, Iran. Emails: r.farahmandfar@sanru.ac.ir; m.asnaashari@yahoo.com

Abstract

Kinetic analysis of gallic acid and methyl gallate in purified Kilka oil was studied in the concentration, range 200–1600 ppm, during autooxidation in Rancimat test at 60°C. The stabilization factor (F), the oxidation rate ratio (ORR), the activity (A), and the mean rate of antioxidant consumption (W_{inh}) were determined. The scavenging activity of 1,1-diphenyl-2-picrylhydrazyl (DPPH) radicals decreased in the order gallic acid > methyl gallate > α -tocopherol > BHT as observed in Kilka oil. There were no significant differences between the effectiveness and strength and activity of gallic acid and methyl gallate at concentrations 200, 400, and 800 ppm. However, above 800 ppm, the activity of methyl gallate became higher than gallic acid, because methyl gallate despite gallic acid did not participate in any chain propagation reactions.

KEYWORDS

antioxidant activity, gallic acid, kilka oil, kinetic, methyl gallate, rancimat

1 | INTRODUCTION

Long-chain polyunsaturated fatty acids (PUFAs) are identified as healthiness compounds observed in marine oil. The marine fish oils are renowned sources of PUFAs, namely eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), which were health beneficial and potential for preventing diseases such as cancer and immune disorders. Besides, omega-3 polyunsaturated fatty acids of fish oil are essential in brain and muscle development and lower plasma triglyceride levels (Horrocks & Yeo, 1999; Uauy & Valenzuela, 2000). On the other hand, highly unsaturated fish oils are notably prone to oxidation and produce unlikeable rancid smells and off-flavors (Jacobsen, Adler-Nissen, & Meyer, 1999).

In the food industry, some food deterioration, nutritional losses, and unpleasant flavors occur during lipid oxidation. The distasteful changes in edible oils increase when reacting with oxygen. Oxidative rancidity cannot be stopped by reducing the storage temperature because it is a chemical reaction with low activation energy (Asnaashari, Tajik, & Khodaparast, 2015; Farhoosh, Johnny, Asnaashari, Molaahmadibahraseman, & Sharif, 2016). Antioxidants are utilized as food ingredient to prolong the shelf life of oils during frying and storage. Using antioxidants are evaluated by different factors including effectiveness and legislation cost (Asnaashari, Hashemi, Mehr, & Asadi Yousefabad, 2015; Sun, Wang, Chen, & Li, 2011). Both natural and synthetic antioxidants have been developed to preclude rancidity and undesirable flavor in fish oil (Horn, Nielsen, & Jacobsen, 2009). Due to positive effects of natural antioxidants, as a consequence of the different side effects of synthetic antioxidants, the consumer demands have extraordinarily improved (Asnaashari, Asnaashari, Ehtiati, & Farahmandfar, 2015).

Gallic acid as a natural phenolic compound and its esters like methyl, octyl, and propyl gallates are broadly utilized in cosmetic, pharmaceutical, and food manufacturing. Gallic acid is originated in variety of plants comprising strawberries, barberry, mango peels, leaves, and walnut (Asnaashari, Farhoosh, & Sharif, 2014). Gallic acid accomplishes as an effective antioxidant and facilitates to protect our cells contrary to oxidation and preclude heart disease (Farhoosh, Sharif, Asnaashari, Johnny, & Molaahmadibahraseman, 2016;

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. Food Science & Nutrition published by Wiley Periodicals, Inc.

FV_Food Science & Nutrition

Polewski, Kniat, & Slawinska, 2002). Although the antioxidative properties of the phenolic acids were considered in the numerous researches (Eshghi, Asnaashari, Haddad Khodaparast, & Hosseini, 2014; Farahmandfar, Asnaashari, & Sayyad, 2015; Thurmann & Herrmann, 1980; Yanishlieva & Marinova, 1995), there is no kinetic research in the literature about the mechanism of gallic acid and methyl gallate.

The objective of this kinetic study was to obtain fundamental information about antioxidative behavior of gallic acid and methyl gallate at 60°C in the Rancimat test at various concentration levels (200, 400, 800, and 1,600 ppm). Furthermore, the activity and the mechanism of these two antioxidants in the triacylglycerols of Kilka oil have been discussed.

2 | MATERIALS AND METHODS

2.1 | Materials

No added components of crude Kilka oil with distinct physicochemical properties (Table 1) were provided from Parskilka Company. Antioxidants like gallic acid, α -tocopherol, BHT, and methyl gallate were purchased from Merck and Sigma Chemical Companies. The structural formulas of these antioxidants are shown in Figure 1. The solvents and chemicals were as analytical reagent grade and obtained from Sigma and Merck Chemical Companies.

2.2 | Purification of the Kilka oil

Kilka oil (120 g) was purified chromatographically to eliminate natural antioxidants. Three absorbents consist of 60 g aluminum oxide 60 (activated at 200°C for 3 hr, neutral) in bottom of the column, 80 g silica gel (activated at 160°C for 3 hr, 60–200 mesh), and 2 g carbon active on the top were used in glass column (50×5 cm i.d.) sequentially. The collection vessels and chromatographic column were covered by aluminum foil, and the suction (without solvent) draw the oil through the column.

TABLE 1	Physicochemical	characteristics	of the crude	Kilka oil ^a
---------	-----------------	-----------------	--------------	------------------------

Parameter	Amount
Wax content (%)	11.07 ± 0.68
Unsaphonifable matter (%)	2.60 ± 0.22
Sterol content (%)	7.523 ± 0.28
Total tocopherols content (mg α -tocopherol per kg oil)	2.89 ± 4.89
Total phenolic content (mg gallic acid per kg oil)	134.02 ± 6.61
Saponification number (mg KOH per g oil)	53.72 ± 3.54
Peroxide value (meq O ₂ per kg oil)	1.78 ± 0.04
Acid value (mg KOH per g oil)	13.98 ± 0.43
Density (g/cm³)	950.72 ± 0.52
Total polar compounds content (%)	52.73 ± 0.92

^aMean ± SD (standard deviation) of triplicate determinations.

2.3 | Rancimat test

Methyl gallate and gallic acid at different concentration of 200, 400, 800, and 1,600 ppm and also α -tocopherol and BHT (200 ppm) were separately added to purified Kilka oil and then subjected to the 743 Rancimat apparatus from Metrohm at 60°C. Sample size and airflow rate were 3 g and 15 L/s, respectively. Electrodes, measuring vessels, connecting tubes, and glassware were cleaned before the assay.

2.4 | Determination of IP of oil samples

The oxidation stability of Kilka oil samples containing antioxidant was evaluated by the Rancimat instrument. In this technique, the tertiary products of oil oxidation, principally formic acid (C_1), acetic acid (C_2), and propionic acid (C_3), formed under accelerated conditions were identified by continuously measuring the water electrical conductivity over time. The Rancimat test expressed the induction period (IP) as the time before rapid deterioration of the oil occurs.

The antioxidant effects on lipid oxidation of Kilka oil are discussed by kinetic parameters: Stabilization factor (F) points to the probability of chain termination of free radicals, especially peroxide radicals. Stabilization factor (F) has determined by the following formulation:

$$F = IP_{inh} / IP_0$$
(1)

where IP_{inh} and IP_0 are induction period in the presence and absence of antioxidant. The oxidation rate ratio (ORR) is a quantity of strength, which could be calculated by following equation:

$$ORR = W_{inh} / W_0$$
 (2)

where W_{inh} and W_0 are the slopes of the linear initiation stage of lipid oxidation in attendance of an antioxidant and control, respectively.

Activity (A), shown as follows, is a common parameter, which simultaneously indicates the efficacy of an antioxidant in termination of the oxidation process and its capability to decline the oxidation rate during the induction period.



FIGURE 1 Molecular structure of gallic acid, methyl gallate, and α -tocopherol and BHT

$$A = F / ORR$$
 (3)

The mean rate of inhibitor consumption (W_{inH}) determined by the formula:

$$W_{inH} = [inH]_0 / IP_{inh} (Ms^{-1})$$
(4)

where $[inH]_0$ and IP_{inh} are the initial concentration of antioxidant (M) and the length of induction period (s), respectively (Marinova & Yanishlieva, 1992).

2.5 Determination of radical scavenging activity

Total antioxidant ability was determined by DPPH assay. In the methanol solution, the samples were reacted with the stable DPPH' radicals. After 30 min incubation period (25°C, dark situation), the absorbance of the solution was quantified (at 517 nm) by a spectrophotometer. DPPH free radical inhibition (I%) was calculated in the following way (Lima, Fernandes-Ferreira, & Pereira-Wilson, 2006):

$$I\% = (A_{blank} - A_{sample}) / A_{blank} \times 100$$
(5)

where A_{blank} and A_{sample} are the absorption of control and sample solutions, respectively. The IC_{50} value, the concentration of the sample required to inhibit 50% of DPPH radical, was achieved by the linear fitted line of dose-response curve plotting between inhibition and concentrations. Antiradical power (ARP) is inverse of IC_{50} , and a relative ARP is the activity of experimented compounds in relation to the strongest scavenger in which the ARP was stated as 100%.

2.6 Statistical analysis

The obtained data were subjected to analysis of variance by SAS software. Significant differences between means were determined by Duncan's multiple range tests at the 5% probability level (p < .05).

RESULTS AND DISCUSSION 3

3.1 | Scavenging effect on DPPH radicals

DPPH free radical scavenging by antioxidants is due to their hydrogen denoting ability. The dose-response plotting curve between absorbance and concentrations for the free radical scavenging capacity of BHT, α -tocopherol, methyl gallate, and gallic acid by the DPPH coloring test were presented in Figure 2. The absorbance of samples reduced severely with the increase in the concentration of antioxidant to a certain extent and then diminished slightly with further concentrations. At a dosage of 20 mg/ml of compounds, the highest DPPH scavenging ability belonged to gallic acid, which is found to be 97.6% and followed by methyl gallate (85.15%), BHT (29.71%), and α -tocopherol (22.89%). While in 50 mg/ml dosage, the order of DPPH scavenging ability of samples was slightly different because the scavenging ability of α -tocopherol with 56.22% was higher than BHT with 43.79%. Indeed, the various radical



FIGURE 2 Radical scavenging activities of gallic acid and methyl gallate, α -tocopherol, and BHT on DPPH, as measured by changes in absorbance at 517 nm

scavenging observed can be associated with different abilities of individual phenolic acids to react with DPPH to give a stable nonradical product.

Values for half-inhibition concentration (IC₅₀), antiradical power (ARP), and a relative ARP of experimented compounds are showed in Table 2. The antiradical activity drops off in the order gallic acid > methyl gallate > α -tocopherol > BHT. Actually, gallic acid required lower doses than methyl gallate, α -tocopherol, and BHT to reach 50% DPPH' scavenging ability. The high value of scavenging activity for gallic acid was reported by (Karamac, Kosiñska, & Pegg, 2005). In the phenolic acids, hydroxyl groups attached to the benzoic molecule are important for free radical scavenging efficiency. As can be seen in Table 2, gallic acid, with four hydroxyl groups, was the strongest antioxidant. Researchers showed that methylation of OH group diminished ARP value in methyl gallate (Kikuzaki, Hisamoto, Hirose, Akiyama, & Taniguchi, 2002).

3.2 | Kinetic study of gallic acid and methyl gallate in triacylglycerols of Kilka oil

Fatty acids composition of the Kilka fish oil is shown in Table 3. This oil was predominantly constituted of oleic, palmitic, palmitoleic, linoleic, myristic, eicosapentaenoic, and docosahexaenoic acids. Among the fatty acids, the maximum amount of the polyunsaturated, monounsaturated, and saturated fatty acids were linoleic acid (8.16%), oleic acid (27.51%), and palmitic acid (17.31%), respectively. The fatty acids composition of Kilka oil showed that the amount of EPA (6.35%) and DHA (5.89%) was marvelous. The fatty acids composition of the Kilka oil used in this study was by data reported in the literature (Frankel, Satué-Gracia, Meyer, & German, 2002). Furthermore, (EPA + DHA)/C16:0 ratio was 0.71 and the total content of PUFA and ω -3 fatty acids were 21.77 and 13.40%, respectively.

TABLE 2 Antiradical activities of selected phenolic acids against

 DPPH^a

Compound	IC ₅₀	ARP	Relative ARP
Gallic acid	5.006	0.2	100
Methyl gallate	7	0.142	71.514
α -Tocopherol	45.35	0.022	11.039
BHT	50.003	0.019	10.011

^aMean ± SD (standard deviation) of triplicate determinations.

TABLE 3 Fatty acid composition of purified Kilka oil

Fatty acid	Percentage (%)
C14:0	6.22 ± 0.05
C16:0	17.31 ± 0.01
C16:1	13.23 ± 0.07
C17:0	1.89 ± 0.06
C18:0	3.23 ± 0.04
C18:1	27.51 ± 0.06
C18:2	8.16 ± 0.15
C18:3	1.17 ± 0.09
C20:0	1.16 ± 0.06
C20:4	0.21 ± 0.03
C20:5 (EPA)	6.35 ± 0.05
C22:6 (DHA)	5.89 ± 0.03
SFA	29.80 ± 0.12
MUFA	40.74 ± 0.01
ω3	13.40 ± 0.11
PUFA	21.77 ± 0.06
PUFA/SFA	0.73 ± 0.01
(EPA + DHA)/C16:0	0.71 ± 0.002
IV [g l ₂ /100 g oil]	114.99 ± 1.10

The remarkable content of these fatty acids make Kilka oil as an important functional food. However, due to its high susceptibility to oxidation, antioxidants need to be added.

Mechanism of lipid oxidation in a noninhibited and inhibited kinetic regime (a sufficiently high oxygen concentration) is presented in Figure 3. Adding an antioxidant into the lipid system causes modification in the process kinetics. In this study, the effect of the gallic acid, methyl gallate, α -tocopherol, and BHT in triacylglycerol's of Kilka oil relies on their molecules and their result radicals in a series of reactions (Scheme 1). In the following schemes, LH is the oxidizing lipid substrate, InH is the inhibitor, in' is the radical of antioxidant, and LOO' is the peroxide radical.

The attained parameters for gallic acid and methyl gallate (200, 400, 800, and 1,600 ppm) and 200 ppm of α -tocopherol and BHT, after processing the kinetic curves, are given in Table 4. It can be seen that as the concentration of antioxidants raised, the efficacy enhanced because of chain termination of free radicals using interaction with peroxide radicals (reaction 7). The strength of antioxidants

0

0.	$2RH + O_2 \rightarrow 2R' + H_2O_2$
1.	$R' + O_2 \rightarrow R00'$
2.	$R00^{\circ} + RH \rightarrow ROOH + R^{\circ}$
3.	ROOH \rightarrow RO' + OH'
3′.	$2\text{ROOH} \rightarrow \text{ROO'} + \text{RO'} + \text{H}_2\text{O}$
3″.	$ROOH + RH \rightarrow RO' + R' + H_2O$
4.	$-R' + R' \rightarrow R - R$
5.	$R' + ROO' \rightarrow ROOR$
6.	ROO' + ROO' \rightarrow products
7.	$ROO' + InH \rightarrow ROOH + In'$
7′.	$In' + ROOH \rightarrow InH + ROO'$
8.	$In' + ROO' \rightarrow In - OOR$
9.	$In' + In' \rightarrow products$
10.	$In' + RH \rightarrow InH + R'$
11.	$InH + ROOH \rightarrow In^{} + RO^{} + H_2$
12.	$InH + O_2 \rightarrow In^- + HOO$
13.	InOOR \rightarrow InO' + RO'
14.	$\ln' + 0_2 \rightarrow \ln 00^{\circ}$

FIGURE 3 Noninhibited (0–6) and inhibited (7–14) oxidation reaction. InH, inhibitor; RH, oxidation lipid substrate; ROO⁻, peroxide radical.

which specifies the chance of antioxidant moieties participating in other reactions, for example 7, 10, 11, 12, and 14, also increased in higher concentrations. Indeed, the possibility of occurring reactions 7-14 relies on the antioxidant structure, temperature, lipid medium, and other conditions of the oxidation process (Marinova & Yanishlieva, 2003). As shown in Table 3, the effectiveness, strength, and activity of gallic acid and methyl gallate were not significantly different, except 1,600 ppm of methyl gallate, which was remarkably higher than gallic acid. The activity (A parameter) of 200 ppm of experimented compounds decreased in the following order: gallic acid ~ methyl gallate > α -tocopherol > BHT. By comparing the order of antioxidants activity in oil with the DPPH radical scavenging activity, it is obvious that methyl gallate was almost as same effective as gallic acid, despite being less polar and radical scavenging activity. Interactions between the methoxy group of methyl gallate and triglyceride molecular results in meaningful antioxidant activity in edible oil. Also, intramolecular interactions between the oxygen atom of the hydroxyl group and the hydrogen atoms of the methylene group of methyl gallate in the meta position of the gallate can increase the hydrogen donor ability and antioxidant activity of methyl gallate (Stöckmann, Schwarz, & Huynh-Ba, 2000).

Figure 4 shows dependency of the stabilization factor (F) on the concentrations of gallic acid and methyl gallate through oxidation of Kilka oil (at 60°C). As can be seen in Figure 4, in lower concentrations, the effectiveness of gallic acid was basically as same as that of methyl gallate. However, at concentrations above $4 \times 1,000$ M methyl gallate were more efficient as compared with gallic acid.

Nonlinear dependence of the F with gallic acid and methyl gallate concentrations is attributed to the participation of their molecules in reactions other than the main reaction of chain termination (7),

TABLE 4 Kinetic parameters characterizing the inhibited lipid oxidation of purified Kilka oil at 60°C in PV₂ = 0 meq/kg, IP₂ = 1.60 hr, W₂ = 68.1 × 10⁻⁹ M/S

	Inhibitor conce	ntration					
Antioxidant	[InH] ^a mg/L	[InH]×1,000(M)	F ^b	ORR ^c	A ^d	W _{inh} (×10 ⁹) (M/s) ^e	W _{InH} (×10 ¹⁰)(M/s) ^f
Gallic acid	200	1.18	11.53	0.38	30.38	23.5	177.32
	400	2.35	19.64	0.34	57.52	21.3	207.3
	800	4.70	26.31	0.27	97.56	16.6	309.5
	1,600	9.41	32.47	0.17	186.5	10.8	502.1
Methyl gallate	200	1.09	9.96	0.37	26.63	23.1	189.1
	400	2.17	16.57	0.30	53.58	19.1	226.9
	800	4.34	26.04	0.28	93.14	17.4	288.8
	1,600	8.69	42.54	0.21	194.6	13.5	353.9
BHT	200	0.91	7.96	0.37	21.45	23.1	197.9
α -Tocopherol	200	0.46	6.98	0.32	24.37	13.2	105.4

^aConcentration of the antioxidant.

^bStabilization factor.

^cOxidative rate ratio.

^dActivity.

^eOxidation rate of inhibited oxidation (M/s).

^fMean rate of inhibitor consumption (M/s).



FIGURE 4 Dependence of gallic acid and methyl gallate stabilization factor F on its concentration [InH] during oxidation of Kilka oil at 60°C

namely reaction (11) and/or (12). In this case, there is a relationship between the mean rate of antioxidant consumption W_{InH} and the antioxidant concentration [InH] (Emanuel, Denisov, & Maizuss, 1965):

$$W_{\ln H} = W_i / f + K_{eff} [\ln H]^n$$
(6)

where W_i is the mean rate of inhibition during the induction period of the inhibited oxidation (M/s), f is the stoichiometric coefficient of inhibition, which determines how many radicals perish in an inhibitor molecule, and K_{eff} is the rate constant of antioxidant consumption in side reaction(s) of chain propagation (Marinova & Yanishlieva, 2003).

The dependence of W_{InH} on [InH] results for gallic acid and methyl gallate at 60°C showed linearity for n = 1 (Figure 5a), which means that these two antioxidants only participate in one of the two side reactions of chain propagation (11 or 12). Indeed, reaction 11 is dependent on the hydroperoxide reactivity. So, the main side reaction in which the antioxidants in inhibited lipid oxidation participate



FIGURE 5 (a) Dependence of the mean rate of consumption W_{inH} of gallic acid and methyl gallate on their concentration [InH] b) Dependence of the antioxidant activity of gallic acid and methyl gallate on the concentrations during the oxidation of Kilka oil at 60°C



FIGURE 6 (a) Dependence of the rate of inhibited oxidation W_{inh} on the concentration of phenolic acids $[InH]^{-1}$ (b) Dependence of the rate of inhibited oxidation W_{inh} on the concentration of phenolic acids $[InH]^{-0.5}$ during oxidation of Kilka oil at 60°C

is reaction 11 (Marinova & Yanishlieva, 1996). From the slope of the dependencies in Figure 5a, the rate constants of this side reaction were 40.256×10^{-7} s⁻¹ and 21.09×10^{-7} s⁻¹ for gallic acid and methyl gallate, respectively. Previous investigations on the antioxidative behavior of some phenolic acids in different lipid systems indicated that K_{eff} depended in the lipid substrate. The parameter W_i/*f* shows the participation of the antioxidant in the initiation reactions. W_i/*f* as determined from Figure 5a by extrapolation to zero concentration of gallic acid (121/55 M/s) and methyl gallate (178.99 M/s), which indicated that methyl gallate participates in chain initiation more than gallic acid during the oxidation. This fact may be one of the reasons for the slightly lower effectiveness and strength of methyl gallate than gallic acid.

Figure 5b clarifies the changes of antioxidant activity with increasing gallic acid and methyl gallate concentrations in the oxidation process of purified Kilka oil at 60°C. As can be seen in Figure 5b, the activities of gallic acid and methyl gallate are practically the same in the concentration interval 0.001–0.006 M. However, above this concentration the activity of methyl gallate is slightly higher than gallic acid.

In order to evaluate the participation of antioxidant radical ln on chain propagation reactions, the following relations developed. The previous research showed that if In' does not participate in chain propagation, this relationship is valid (Marinova & Yanishlieva, 1996).

$$W_{inh} \sim [\ln H]^{-1}$$
(7)

While if In participates in one of the chain propagation (reaction 7, 10, and 14), this relationship is observed.

$$W_{inb} \sim [\ln H]^{-0.5}$$
 (8)

Processing of the results obtained from the dependence of the rate of inhibited oxidation W_{inh} on the concentrations of gallic acid and methyl gallate (7) showed the relationship was valid (Figure 6a). This means that the radicals of methyl gallate did not participate in any chain propagation. However, gallic acid indicated no linear dependence on both $[InH]^{-1}$ and $[InH]^{-0.5}$, which showed that the radicals of gallic acid were included in more than one reaction of chain propagation (Figure 6b). As a consequence, the stability of oil including gallic acid at 1,600 ppm is lower than oil contained methyl gallate. Moreover, based on (7) relationship for both antioxidants showed a good linear relationship at concentration lower than 800 ppm. It means that lower than 800 ppm, radicals of gallic acid and methyl gallate did not participate in chain propagation reactions. But at 1,600 ppm the mechanism became different.

4 | CONCLUSIONS

The results of the present study indicated that methylation of benzoic acid could change the kinetic mechanism of oxidation in Kilka oil. According to the kinetic parameters obtained for gallic acid and methyl gallate, there was observed no significant differences between the activity, effectiveness, and strength of these two antioxidants at 200–800 ppm concentrations, whereas at 1,600 ppm, methyl gallate acts better than gallic acid due to no participation in any propagation chain reactions. Moreover, comparison of the activity of gallic acid and methyl gallate with α -tocopherol and synthetic antioxidant (BHT) indicated that these two antioxidants have better potential to inhibit lipid oxidation in Kilka oil. So, they can be used as suitable alternatives to synthetic antioxidants.

ACKNOWLEDGMENTS

We are grateful to Ferdowsi University of Mashhad (FUM) and Sari Agricultural Sciences and Natural Resources University (SANRU).

CONFLICT OF INTEREST

The authors have declared no conflict of interest.

ETHICAL APPROVAL

This study does not involve any human or animal testing.

ORCID

Reza Farahmandfar D https://orcid.org/0000-0002-0037-4139

REFERENCES

- Asnaashari, E., Asnaashari, M., Ehtiati, A., & Farahmandfar, R. (2015). Comparison of adaptive neuro-fuzzy inference system and artificial neural networks (MLP and RBF) for estimation of oxidation parameters of soybean oil added with curcumin. *Journal of Food Measurement* and Characterization, 9(2), 215–224. https://doi.org/10.1007/ s11694-015-9226-7
- Asnaashari, M., Farhoosh, R., & Sharif, A. (2014). Antioxidant activity of gallic acid and methyl gallate in triacylglycerols of Kilka fish oil and its oil-in-water emulsion. *Food Chemistry*, 159, 439–444. https://doi. org/10.1016/j.foodchem.2014.03.038
- Asnaashari, M., Hashemi, S. M. B., Mehr, H. M., & Asadi Yousefabad, S. H. (2015). Kolkhoung (Pistacia khinjuk) hull oil and kernel oil as antioxidative vegetable oils with high oxidative stability and nutritional value. Food Technology and Biotechnology, 53(1), 81–86. https://doi. org/10.17113/ftb.53.01.15.3719
- Asnaashari, M., Tajik, R., & Khodaparast, M. H. H. (2015). Antioxidant activity of raspberry (Rubus fruticosus) leaves extract and its effect on oxidative stability of sunflower oil. *Journal of Food Science* and Technology, 52(8), 5180–5187. https://doi.org/10.1007/ s13197-014-1564-7
- Emanuel, N., Denisov, E., & Maizuss, Z. (1965). Chain radical oxidation of hydrocarbons in liquid phase. Moscow, Russia: Nauka.
- Eshghi, N., Asnaashari, M., Haddad Khodaparast, M. H., & Hosseini, F. (2014). Evaluating the potential of natural curcumin for oxidative stability of soybean oil. *Natural Product Research*, *28*(17), 1375–1378. https://doi.org/10.1080/14786419.2014.901319
- Farahmandfar, R., Asnaashari, M., & Sayyad, R. (2015). Comparison antioxidant activity of Tarom Mahali rice bran extracted from different extraction methods and its effect on canola oil stabilization. *Journal* of Food Science and Technology, 52(10), 6385–6394. https://doi. org/10.1007/s13197-014-1702-2
- Farhoosh, R., Johnny, S., Asnaashari, M., Molaahmadibahraseman, N., & Sharif, A. (2016). Structure-antioxidant activity relationships of ohydroxyl, o-methoxy, and alkyl ester derivatives of p-hydroxybenzoic acid. *Food Chemistry*, 194, 128–134. https://doi.org/10.1016/j. foodchem.2015.08.003
- Farhoosh, R., Sharif, A., Asnaashari, M., Johnny, S., & Molaahmadibahraseman, N. (2016). Temperature-dependent mechanism of antioxidant activity of o-hydroxyl, o-methoxy, and alkyl ester derivatives of p-hydroxybenzoic acid in fish oil. Journal of the American Oil Chemists' Society, 1–13. https://doi.org/10.1007/ s11746-016-2790-0
- Frankel, E. N., Satué-Gracia, T., Meyer, A. S., & German, J. B. (2002). Oxidative stability of fish and algae oils containing long-chain polyunsaturated fatty acids in bulk and in oil-in-water emulsions. *Journal* of Agricultural and Food Chemistry, 50(7), 2094–2099. https://doi. org/10.1021/jf0111458
- Horn, A. F., Nielsen, N. S., & Jacobsen, C. (2009). Additions of caffeic acid, ascorbyl palmitate or γ-tocopherol to fish oil-enriched energy bars affect lipid oxidation differently. *Food Chemistry*, 112(2), 412– 420. https://doi.org/10.1016/j.foodchem.2008.05.094
- Horrocks, L. A., & Yeo, Y. K. (1999). Health benefits of docosahexaenoic acid (DHA). *Pharmacological Research*, 40(3), 211–225. https://doi. org/10.1006/phrs.1999.0495

- Jacobsen, C., Adler-Nissen, J., & Meyer, A. S. (1999). Effect of ascorbic acid on iron release from the emulsifier interface and on the oxidative flavor deterioration in fish oil enriched mayonnaise. *Journal* of Agricultural and Food Chemistry, 47(12), 4917–4926. https://doi. org/10.1021/jf990241u
- Karamac, M., Kosiňska, A., & Pegg, R. B. (2005). Comparison of radicalscavenging activities for selected phenolic acids. Polish Journal of Food and Nutrition Sciences, 14(2), 165–170.
- Kikuzaki, H., Hisamoto, M., Hirose, K., Akiyama, K., & Taniguchi, H. (2002). Antioxidant properties of ferulic acid and its related compounds. *Journal of Agricultural and Food Chemistry*, 50(7), 2161–2168. https://doi.org/10.1021/jf011348w
- Lima, C. F., Fernandes-Ferreira, M., & Pereira-Wilson, C. (2006). Phenolic compounds protect HepG2 cells from oxidative damage: Relevance of glutathione levels. *Life Sciences*, 79(21), 2056–2068. https://doi. org/10.1016/j.lfs.2006.06.042
- Marinova, E. M., & Yanishlieva, N. V. (1992). Inhibited oxidation of lipids II: Comparison of the antioxidative properties of some hydroxy derivatives of benzoic and cinnamic acids. *Fett Wissenschaft Technologie/ Fat Science Technology*, 94(11), 428–432. https://doi.org/10.1002/ lipi.19920941110
- Marinova, E. M., & Yanishlieva, N. V. (1996). Antioxidative activity of phenolic acids on triacylglycerols and fatty acid methyl esters from olive oil. Food Chemistry, 56(2), 139-145. https://doi. org/10.1016/0308-8146(95)00147-6
- Marinova, E. M., & Yanishlieva, N. V. (2003). Antioxidant activity and mechanism of action of some phenolic acids at ambient and high temperatures. *Food Chemistry*, 81(2), 189–197. https://doi.org/10.1016/ S0308-8146(02)00411-9
- Polewski, K., Kniat, S., & Slawinska, D. (2002). Gallic acid, a natural antioxidant, in aqueous and micellar environment: Spectroscopic studies. Current Topics in Biophysics, 26, 217–227.
- Stöckmann, H., Schwarz, K., & Huynh-Ba, T. (2000). The influence of various emulsifiers on the partitioning and antioxidant activity of hydroxybenzoic acids and their derivatives in oil-in-water emulsions. *Journal of the American Oil Chemists' Society*, 77(5), 535–542. https:// doi.org/10.1007/s11746-000-0085-6
- Sun, Y.-E., Wang, W.-D., Chen, H.-W., & Li, C. (2011). Autoxidation of unsaturated lipids in food emulsion. *Critical Reviews in Food Science* and Nutrition, 51(5), 453–466. https://doi.org/10.1080/1040839100 3672086
- Thurmann, I., & Herrmann, K. (1980). Uber die antioxidative Wirkung von Hydroxizimtsauren und Hydroxibenzoesauren. Deutsche Lebensmittel-Rundschau.
- Uauy, R., & Valenzuela, A. (2000). Marine oils: The health benefits of n-3 fatty acids. Nutrition, 16(7), 680–684. https://doi.org/10.1016/ S0899-9007(00)00326-9
- Yanishlieva, N., & Marinova, E. M. (1995). Effects of antioxidants on the stability of triacylglycerols and methyl esters of fatty acids of sunflower oil. *Food Chemistry*, 54(4), 377-382. https://doi. org/10.1016/0308-8146(95)00061-M

How to cite this article: Asnaashari M, Farhoosh R, Farahmandfar R. Preservation of gallic acid and methyl gallate on purified Kilka fish oil oxidation by Rancimat. *Food Sci Nutr.*

2019;7:4007-4013. https://doi.org/10.1002/fsn3.1264