






REVIEW ARTICLE

Gut microbiota and synbiotic foods: Unveiling the relationship in COVID-19 perspective

Noor Akram¹ | Farhan Saeed²  | Muhammad Afzaal² | Yasir Abbas Shah²  |
Aiza Qamar³ | Zargham Faisal⁴ | Samia Ghani⁵ | Huda Ateeq²  |
Muhammad Nadeem Akhtar⁶  | Tabassum Tufail⁶  | Muzzamal Hussain² |
Aasma Asghar² | Ammara Rasheed¹ | Entessar Al Jbawi⁷

¹Department of Food and Nutrition,
Government College University,
Faisalabad, Pakistan

²Department of Food Science,
Government College University,
Faisalabad, Pakistan

³Department of Nutrition and Health
Promotion, University of Home Economics
Lahore, Lahore, Pakistan

⁴Institute of Food Science and Nutrition,
Bahauddin Zakariya University Multan,
Multan, Pakistan

⁵Faculty of Pharmaceutical Sciences,
Government College University
Faisalabad, Punjab, Pakistan

⁶University Institute of Diet and
Nutritional Sciences, The University of
Lahore, Lahore, Pakistan

⁷Agricultural Extension Directorate,
MAAR, Damascus, Syria

Correspondence

Muhammad Afzaal, Department of Food
Science, Government College University,
Faisalabad, Pakistan.

Email: muhammadafzaal@gcuf.edu.pk

Entessar Al Jbawi, Agricultural Extension
Directorate, MAAR, Damascus, Syria.
Email: dr.entessara@nmail.sy

Abstract

The Coronavirus disease 2019 (COVID-19) has spread across the globe and is causing widespread disaster. The impact of gut microbiota on lung disease has been widely documented. Diet, environment, and genetics all play a role in shaping the gut microbiota, which can influence the immune system. Improving the gut microbiota profile through customized diet, nutrition, and supplementation has been shown to boost immunity, which could be one of the preventative methods for reducing the impact of various diseases. Poor nutritional status is frequently linked to inflammation and oxidative stress, both of which can affect the immune system. This review emphasizes the necessity of maintaining an adequate level of important nutrients to effectively minimize inflammation and oxidative stress, moreover to strengthen the immune system during the COVID-19 severity. Furthermore, the purpose of this review is to present information and viewpoints on the use of probiotics, prebiotics, and synbiotics as adjuvants for microbiota modification and its effects on COVID-19 prevention and treatment.

KEYWORDS

COVID-19, gut microbiota, immune system, nutrition, probiotics

1 | INTRODUCTION

The novel coronavirus has been causing a global health disaster since December 2019. COVID-19 is an acute pulmonary health disease caused by "Severe Acute Respiratory Syndrome coronavirus-2" (SARS-Cov-II virus). It has been reported that this virus had infected more than 210 million people and 4.4 million have died globally.

Representative findings include high fever, coughing, sneezing, pneumonia, and loss of taste and smell (Huang et al., 2020). Moreover, 18% of COVID-19 patients often suffer from gastrointestinal problems like diarrhea and vomiting (Cheung et al., 2020).

Some medical conditions and factors like age and sex, and other bad lifestyle factors such as smoking, exert an influence on the host metabolic status which determines the clinical severity of COVID-19

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Food Science & Nutrition* published by Wiley Periodicals LLC.

(Kilgore, 2020). Some prevailing diseases come into view that can make worse the COVID-19 infection, like hypertension, diabetes (type 2), cardiovascular diseases (CVD), cancers, and smoking, all of these factors regulate the illness course, progression, and outcome of COVID-19 (Bae et al., 2021). The most diverse ecological community of microorganisms in the human body is "Gut microbiota." It is another important factor that plays a condemnatory role during pulmonary infection. The human gastrointestinal tract possesses millions of microorganisms, and alterations in the composition of gut microbiota due to different factors is termed "dysbiosis," which has been interlinked with various human diseases. Some studies have used metagenomics sequencing proposed data regarding the clinical findings of COVID-19 patients; the microbial diversity is decreased as compared to pathogens in fecal samples (Yeoh et al., 2021). Moreover, various beneficial microbes like *Eubacterium rectale* and *Faecalibacterium prausnitzii* have been reported to drain in COVID-19 patient's fecal samples. However, opportunistic pathogens such as *Collinsella* species and *Morganella morganii* were exclusively high in fecal samples of SARS-Cov-II patients (Zuo et al., 2020). These studies have concluded that with an increase in pathogenic microorganisms is intently connected with sickness seriousness in SARS-Cov-II infection.

Foods having essential nutritional compounds have been very effective in the severity of SARS-Cov-II. Consumption of nutrients with antioxidant and antiinflammatory potential is beneficial for COVID-19 patients as imprudent production of proinflammatory cytokines and acute inflammation takes place in SARS-Cov-II infection. Moreover, plant-based phenols, omega-3 fatty acids, zinc, vitamins, and polysaccharides are very major dietary nutrients in SARS-Cov-II infection (Iddir et al., 2020). Contrarily, acute changes occur in gut microbiota due to stress, injury, and inflammation, however, diet along with environmental factors can regulate the diversity of gut microbiota (Voigt et al., 2014). The gut microbiota has been demonstrated to be affected by a variety of dietary components; intake of proteins from peas and whey enhances gut beneficial bacteria like *Lactobacillus* and *Bifidobacterium*, on the other hand, whey has been reported to decrease *Bacteroides fragilis* and *Clostridium perfringens* pathogens in the gut (Earley et al., 2015). Essentially, it was noticed that diet low in fat leads to more fecal overflow of *Bifidobacterium*, while a diet high in saturated fat increases the abundance of *Faecalibacterium prausnitzii* (Dominika et al., 2011).

Moreover, a fiber-rich diet encourages the development of healthy microbiota (Le Chatelier et al., 2013). Dietary fibers are considered as microbiota's most accessible carbohydrates which improve intestinal health and also provide energy to the host. However, prebiotics such as polydextrose, maize fiber, and insulin have been reported to boost gut diversity, immunity, digestion, and health, particularly in elderly people (Bouhnik et al., 2007). Gut microbiota and host immunity are correlated. Diet is again a predominant factor that regulates the host metabolism and elaborates the diversity and functions of gut microbiota (Hahn et al., 2017). Disturbances in nutritional status cause impairment of immune responses (De Santis et al., 2015). It is important to understand how the immune system

responds to viral vaccines and how the efficacy of the vaccine can be increased. The efficacy of the vaccine is influenced by various factors like good nutrition, diet, sex, age, genetics, and immunity; these factors have great variation among low-, middle-, and high-income settings (Valdez et al., 2014). Poor immune responsiveness to vaccination has been shown in low-income countries and various factors are associated with this poor response status, such as genetics, beneficial microbiomes, environmental factors, and interlinked pathogens (Parker et al., 2014; Pulendran, 2014). Early-life environmental factors that influence the microbiome such as the delivery method of birth, diet, and hygienic conditions can affect vaccine efficacy (Guaraldi & Salvatori, 2012). Moreover, micronutrient deficiencies are correlated with undernourishment and less diverse microbiota; particularly, in low- to middle-income countries (LMICs), children are affected by vitamin A deficiency (VAD; Wirth et al., 2017). However, VAD remarkably affects vaccine efficacy and mucosal immunity (Hall et al., 2011).

Dietary components like dietary fiber can serve as prebiotics to enhance the abundance of probiotics in the gut. Moreover, plant-based fibers increase the profusion of probiotics such as *Bifidobacterium* and *Lactobacillus*, while decreasing pathogenic bacteria like *Clostridium* (Iddir et al., 2020). The gut produces SCFAs such as propionate, acetate, and butyrate when certain prebiotic dietary fibers are degraded by gut microbes. SCFAs are major immunomodulatory metabolites that can enhance the responsive activities of CD8+ T-cells and B-cells in the immune system (Alameddine et al., 2019; Trompette et al., 2018). Evidence from recent studies has proved that lung infections are severely affected by less food consumption ultimately leading to alteration in the gut microbiome, as the severity of viral infections is correlated with food intake and gut microbiota diversity. It is needed to understand that gut microbiota is very essential in maintaining good health and lessening the severity of disease or viral infections. The gut microbiomes generate metabolites that can systematically change immune responses. The gut microbiota diversity is directly correlated with the host immune system (Groves et al., 2020).

Several bacteria are found in the human respiratory system, and internalization of respiratory viruses occurs mostly through this route. Probiotics are another friendly group of bacteria that, when swallowed or supplied in a certain concentration, have a good impact on human health. Probiotic strains like *Bifidobacterium* and *Lactobacillus* can trap viruses and interfere with their binding to host cell receptors, which is advantageous to the host's health (Kanauchi et al., 2018). Results from various studies have shown that probiotics may be useful in the treatment of viral infections such as COVID-19, viral infections, pulmonary tract infections, cystic fibrosis, chronic inflammatory diseases, PCOs, weight loss, and improving lactation in women (Alberca et al., 2021; Bajinka et al., 2022; Robichaud et al., 2021). Consumption of probiotic-rich meals or supplements has been proven to impact immune function by modifying the microbiota's endogenous metabolic activity (Galdeano et al., 2019). Moreover, the gut microbiota is a complex collection of microorganisms that colonize the mucosal surfaces of the gut. These bacteria

are considered key elements in health due to their significant metabolites, immune system control, and defense of the body against infections (Human Microbiome Project Consortium, 2012). The effects of prebiotics and probiotics on human health are shown in Figure 1.

2 | MAINTAINING A HEALTHY IMMUNE SYSTEM DURING COVID-19 CRISES

World Health Organization had reported COVID-19 as a health emergency. Challenges emerge to resist the severity of infection ideally and enhance immunity. The role of food as a natural source of immunomodulation cannot be overlooked throughout the vaccination campaign. It has been reported that nutrients like probiotics, prebiotics, folic acid, zinc, and vitamins C, D, E, and A are prime modulators for a healthy immune system (Saeed et al., 2016). As an effective immune response depends upon a balanced diet and accurate nourishment of gut microbiota, it is reported that adequate protein intake is essential for antibody production. Micronutrient deficiencies are associated with high-risk infections and the severity of diseases, particularly vitamin A and zinc (Iddir et al., 2020). However, oxidative stress and inflammation extremely regulate the responses of the immune system (Lauridsen et al., 2019). Oxidative stress and inflammation have been shown a great impact on the immune system. Moreover, vitamins like A, D, and C along with some plant-based proteins like polyphenols, carotenoids, and flavonoids have antiinflammatory and antioxidant potential; these dietary nutrients are vital for a strong and healthy immune system (Iddir et al., 2020). Furthermore, vaccination to combat COVID-19 is now available, and the relevance of natural forms of inhibition and treatment cannot be overlooked. In this regard, food and dietary habits are important factors in determining overall health and resistance (Lindgren et al., 2018).

2.1 | Proteins

According to the Institute of Medicine, the USA recommended daily allowance (RDA) for protein 0.8/kg body weight. Increased risk of infection has been recognized in low-income countries as they are protein malnourished due to low protein availability (Rodríguez et al., 2011). Immunoglobulins and lymphoid tissues associated with the gut play important role in defense against infection; these components are interlinked with protein availability (Amaral et al., 2006). Moreover, studies have reported that very low protein intake decreases the antibodies abundance and increases the risk of viral infections (Taylor et al., 2013). Protein intake of high biological value and essential amino acids may lower inflammation during infection and improve satiety (Chungchunlam et al., 2017). High-quality proteins have proven primary component of the antiinflammatory diet (O'Keefe et al., 2008). Branched-chain amino acids enhance the gut barrier by increasing the immunoglobulin level in the intestine (Ren et al., 2015).

2.2 | Lipids

The number and type of lipids in the diet have an impact on the composition and function of the gut microbiota. Therefore, dietary lipids can have an impact on host physiology via interacting with the microbiota in the gut. Lipids have an impact on the gut microbiota both as substrates for bacterial metabolic processes and by suppressing bacterial development (Schoeler & Caesar, 2019). Recent studies indicate that cellular metabolism, including lipid metabolism, affects macrophage activity. Lipids provide energy to macrophages and serve as the building blocks for bioactive lipids and cellular membranes. During macrophage activation, lipids also control signaling and gene regulation (Yan & Horng, 2020).

However, antiinflammatory effects contribute to a healthy immune system. Fatty acids can remarkably change immune responses.




FIGURE 1 Adjuvant effect of prebiotics and probiotics on human health

Particularly, fatty acids (FAs) regulate the immune cell functioning such as macrophages, neutrophils, innate lymphoid cells, T-cells, and B-cells (Radzikowska et al., 2019). Two essential fatty acids omega-3 and omega-6 cannot be produced by the human body. Both these FAs are the most potent antiinflammatory nutrients and must be taken through diet (Innes & Calder, 2018; Serhan & Levy, 2018).

2.3 | Carbohydrates

The role of gut microbiota in the immune system is vital and its modulation by adequate nutrition affects the severity of viral infection, but unfortunately, studies regarding, diet, gut, and infection are few (Dubourg et al., 2019). Dietary fibers are complex carbohydrates that play a major role in inflammation (Galland, 2010). Various kinds of dietary fiber are not the sole forces for stomach or gut well-being, but they can help in the production of SCFA that nourishes gut microbiota species. SCFAs level is regulated by diet and plays an important role in immunomodulatory function (Parada Venegas et al., 2019). Moreover, foods with a high glycemic index such as processed foods have been reported to cause increased production of free radicals, which ultimately lead to oxidative stress and inflammation (O'Keefe et al., 2008). Contrarily, the consumption of dietary fiber is linked to a decreased mortality rate because of respiratory and infectious diseases in adult Americans (Park et al., 2011).

2.4 | Minor nutritional components

Vitamin A is associated with the development of solid bodily fluid layers or mucous, particularly, for the respiratory and the digestive tract needed for mucin secretion and upgrading antigen levels (Watchorn et al., 2022). Globally, vitamin A is the most common micronutrient deficiency specifically in low- and middle-income countries where meat intake is low (Ross, 2010). Primarily, a deficiency of vitamin A is linked with an increased probability of infection (Huang et al., 2018). Active forms of vitamin A are retinal, retinol, and retinoic acid, however, retinoic acid among them plays a major role in the maturation, differentiation, regulation, and functioning of the innate immune system (Shrestha et al., 2017). Vitamin A boosts tissue antigen immunity by boosting mucin production in both the gut and respiratory system (Huang et al., 2018). As a result, it is predictable that vitamin A deficiency has been linked to impaired vision. Neutrophils, macrophages, and B- and T-cells all play a role in the maintenance of a healthy immune system (Singh et al., 2001). Moreover, it increases the levels of immunoglobulin A, G, and M in the blood and improves the functioning of the innate immune system (Huang et al., 2020). Furthermore, people with low vitamin A levels have morphological alterations in their epithelial and parenchyma cells lungs, which increases the risk of lung illness and dysfunction (Timoneda et al., 2018). This is critically significant in terms of COVID-19's impact on pulmonary function (Siddiqi & Mehra, 2020).

Vitamin D is obtained through eggs, fortified milk, fish, and mushrooms in the diet. It may also be produced under the skin in the presence of UV light from cholesterol (Mosekilde, 2005). When it comes to the immune system, vitamin D has been shown to modulate both the innate and adaptive immune systems (Childs et al., 2020). The active form of vitamin D is calcitriol (1,25 dihydroxy vitamin D), synthesized by the hydroxylation mechanism in the liver and kidney that is involved in calcium homeostasis, and ultimately, enhances bone health; however, it has also been proven to influence immune system (Mosekilde, 2005). The function of the T-cell is strongly connected to vitamin D status (von Essen et al., 2010). Toll-like receptors (TLR) in the innate system let macrophages recognize lipopolysaccharide (LPS), which acts as a replacement for bacterial immunity (Kaminogawa & Nanno, 2004).

A meta-analysis had shown that vitamin D therapy remarkably improves the severity of chronic obstructive pulmonary disease (COPD) (Li et al., 2020). Another analysis revealed that there was a lower incidence of COVID-19 and influenza infections along with death (Grant et al., 2020). A recent retrospective analysis of confirmed cases of SARS-CoV-2 infection had shown that low vitamin D levels were significantly linked to a higher risk of mortality, particularly, in geriatric and male individuals with prior illnesses (Raharusun et al., 2020).

High intake of ascorbic acid or vitamin C has been linked to lower C-reactive protein (Wannamethee et al., 2006). Vitamin C is essential for immune system function, particularly immune cell activity; it is acknowledged as a traditional antioxidant that quenches free radicals immediately and prevents proinflammatory diseases. This vitamin boosts phagocytosis, boosts oxidant production, boosts microbial death, and speeds up neutrophil transit to the infection site. Its role as a cofactor for various gene regulatory and biosynthetic enzymes is thought to have a role in immunological modulation (Carr & Maggini, 2017). Vitamin C has been demonstrated to increase neutrophil migration to the infection site, triggering the production of reactive oxygen species (ROS) and phagocytosis (Carr & Maggini, 2017; Carr & Melcher, 2017). A clinical trial among 50 COVID cases was performed, in which high-dose vitamin C intravenous intervention was given and outcomes have shown positive changes in the oxygenation index of COVID patients (Cheng, 2020). Adequate levels of vitamin C have been proposed as a therapy treatment for those suffering from the common cold and pneumonia (Hemilä, 2017).

Zinc is a trace element that the human body cannot produce or store, thus it must be obtained via food (Stefanidou et al., 2006). Zinc inadequacy is a severe issue in public health that affects people globally (Wessells & Brown, 2012). Individuals with zinc deficiency are more prone to viral infections (Read et al., 2019). Zinc is also thought to be essential for T-cell receptors that need the intracellular binding of tyrosine kinase, which is necessary for their growth and survival as well as activation of T-lymphocyte activation (Wintergerst et al., 2006). Zinc plays a crucial role in cytokine release and triggering CD8+ T-cell proliferation, as well as immune cell development and differentiation. It also helps to modify cytokine release and

induce CD8+ T-cell proliferation (Wintergerst et al., 2007). Zinc enhances the production of macrophages and phagocytosis, elevates cytokine levels, promotes B- and T-cells growth, and also inhibits free radicals-induced injury (Wessels et al., 2017). In a recent study, the impact of poor zinc status on the elderly and its link to pneumonia were emphasized (Barnett et al., 2010). Essential nutrients in the therapy of COVID-19 and other infections are presented in Table 1.

3 | VACCINE EFFICACY OF COVID-19

The immunological response to the SARS-CoV-2 vaccine and the vaccine's side effects are likely to be influenced by the gut microbiome. According to preliminary results of an ongoing clinical investigation, significant changes were observed in subjects who took a probiotic formula for 2 months and had greater blood SARS-CoV-2 IgG antibody levels and lower proinflammatory cytokines. Large public-based research in the United States enlisted more than 10,000 people to figure out the link between COVID-19 vaccination effectiveness and gut microbiota. The potential role of gut microbiota in protection against SARS-CoV-2 infection is close to the bottom (Lau & Yu, 2022). In children living in low-income and middle-income countries, undernourishment and repetitive gastrointestinal infections are associated with the failure of oral vaccines. Mechanistic studies of the relationship among the microbiome, host genetics, and viral infections are needed to understand how immune responses to viral vaccines can be modulated (Vlasova et al., 2019).

It is currently unclear to what extent SARS-CoV-2 vaccinations provide universal protection, and scientists are working to find out. Researchers from all around the world are seeking to find the optimal quantity of self-antigens or immunological markers. The efficacy of the COVID-19 vaccination is tightly linked. The increase in the

number of instances caused by the delta variation throughout the world has encouraged several governments to think about booster vaccinations for at-risk populations regardless of unsubstantiated arguments (Callaway, 2021). In 2019, the gut microbiome was shown to alter vaccine effectiveness after healthy participants taking antibiotics before the H1N1 influenza vaccination showed a substantial reduction in antibody response (Hagan et al., 2019).


4 | GUT-LUNG AXIS AND POTENTIAL ROLE OF THE GUT IN IMMUNITY

When the virus (SARS-CoV-2) enters host cells, the interface between innate and adaptive immunity is rapidly activated (Chowdhury et al., 2020). Trillions of microorganisms live in the human gut microbiota, including viruses, archaea, bacteria, and fungi (Gill et al., 2006). *Bacteroidetes*, *Proteobacteria*, *Actinobacteria*, and *Firmicutes* are the major four phyla that are dominant in the gut of healthy individuals (Villanueva-Millán et al., 2015). There is a gut-lung axis that facilitates intense communication between the two organs, including microbial and immunological interactions (Enaud et al., 2020). The host provides the bacteria home and food and the microorganisms support the host by providing protective immunity against viral infections and also regulating various physiological activities such as nutritional digestion. A variety of disorders is considered with less diverse gut microbiota (Mosca et al., 2016). Moreover, premature birth, cesarean section, poor breastfeeding, and antibiotic therapy, alone or in combination, are all linked to inefficient microbiota development. During infancy, immunological problems or inflammatory illnesses are common. Changes in the gut microbiota, often known as "gut dysbiosis," have been linked to a variety of diseases and illnesses, including type 2 diabetes, IBD, cardiovascular disease, and

TABLE 1 Essential nutrients in the therapy of Covid-19 and other infections

Nutrient	Study	Outcomes	References
Vitamin D	In Indonesia, two cohort studies of 780 patients with proven SARS-CoV-2 infection were performed. COVID-19 mortality patterns with a special focus on vitamin D deficiency and related variables were investigated in this study	COVID-19 mortality is highly linked to vitamin D deficiency	Raharusun et al. (2020)
Folic acid	The study proposed that pregnant women have a less chance of acquiring infection, particularly SARS-Cov-II, because of FA supplementation	Pregnant women with low folate RBC concentrations who do not take FA supplements had a higher chance of hospitalization. FA supplementation may protect against SARS-CoV-2, according to this research	Acosta-Elias and Espinosa-Tanguma (2020)
Ascorbic Acid	A preliminary study reveals the intravenous effects of vitamin C in patients with sepsis and ARDS, as well as molecular indicators of inflammation and vascular damage. In this study, vitamin C intravenous infusions were given to patients randomly every 6 h for 96 h	During infection, ascorbic acid (AA) alters ROS production, consequently, prevents cell death	Truwit et al. (2019)
Zinc	In vitro study was performed through the inhibition of RNA-dependent RNA polymerase (RdRp) and 3C-like viral proteinase (3CLpro)	Zinc (Zn) inhibits SARS-CoV-2 replication and regulates IL-1 β and TNF- α levels and also modulates cytokine storm in COVID-19	Pormohammad et al. (2021)

FIGURE 2 Sources of prebiotics and probiotics



depression (Dhar & Mohanty, 2020). Several lung disorders, including allergies, asthma, and cystic fibrosis, have been linked to dysbiosis in the gut microbiota (Anand & Mande, 2018). Despite increased food consumption, deficiency in specific nutrients is typically noted in overweight and obese individuals due to an imbalanced diet (Zhang, Liu, et al., 2020). The microbiota may have a key role in the immune response to bacterial infections in the lungs, but the mechanisms behind this need to be researched further (Enaud et al., 2020).

“Gut-lung axis” is a critical cross-talk between the gut microbiota and lungs; gut microbiota is found to influence lung health through this link (Keely et al., 2012). This cross-talk is considered to be bidirectional and endotoxins, and microbial metabolites may influence the lungs causing inflammation that can also affect the gut microbiota (Dumas et al., 2018). Bacteria from the groups *Prevotellaceae* and *Ruminococcaceae*, *Bacteroidaceae*, *Lachnospiraceae*, and *Rikenellaceae* are abundant in the colon (Hall et al., 2017). Numerous pieces of evidence imply the presence of different bacteria in the lungs, similar to the gut microbiota (Bingula et al., 2017). *Bacteroidetes*, *Proteobacteria*, and *Firmicutes* are dominant in the lungs, whereas *Firmicutes* and *Bacteroidetes* dominate in the gut (Zhang, Zhu, et al., 2020). The dysbiosis microbiome may trigger an inflammatory response that the coronavirus can take full advantage of. The dysbiosis microbiome may trigger an inflammatory response that the coronavirus can take full advantage of. When a coronavirus infects the gut, inflammatory substances called cytokines are increased by additional cytokines. The resulting inflammation might set off a “cytokine storm,” a rogue immune response that can do more harm than the virus itself, causing multiorgan impairment (Yang et al., 2020).

Respiratory virus infections are exceedingly prevalent, but little is known about how they affect the makeup and function of the gut microbiota. After a viral lung infection, a study had shown considerable alteration in the gut microbiota. Weight loss during RSV or influenza virus infection was due to reduced food consumption (Groves et al., 2020). This offers the intriguing prospect that new SARS-CoV2 might affect the gut microbiome as well. Alteration in gut microbiota has been linked to respiratory infections in multiple studies (Groves et al., 2020). The lung microbiota becomes abundant in bacteria from the gut in ARDS, most likely due to a hyperpermeable gut that allows germs to pass past the colon wall and into the lungs (Fanos et al., 2020).

Two significant clinical symptoms of COVID-19 are acute respiratory distress syndrome (ARDS) and pneumonia, particularly, in older

individuals and patients with a weak immune system (Lake, 2020). Elderly individuals with a weak immune system have shown substantial negative clinical manifestations. Gut microbiota interactions with the host are various, intricate, and bidirectional. The gut is thought to have a key role in the development and function of adaptive and innate immune systems (Negi et al., 2019). However, elderly adults have less diversified microbiota and beneficial microbes such as *Bifidobacterium* are deficient in the gut (Nagpal et al., 2018). Improving gut microbiota could be one of the prophylactic ways by which the impact of this disease can be minimized in old people and immune-compromised patients (Dhar & Mohanty, 2020). The gut microbiota and immunological homeostasis appear to have a strong correlation. Antimicrobial peptides are produced in the gut when beneficial microbes compete for nutrients and habitat (Moens & Veldhoen, 2012). Furthermore, the derived signals from gut microbiota adjust immune cells for pro- and anti-inflammatory responses to enhance immunity, which may reduce disease susceptibility and progression (Negi et al., 2019). Furthermore, it has been shown that the gut and lungs share a mucosal immune system and that the gut–lung axis links the inflammatory process and immunological responses (Antunes et al., 2020). Different sources of prebiotics and probiotics are shown in Figure 2.

5 | IMPACT OF PROBIOTICS AND SYNBIOTIC ON THE SEVERITY OF COVID-19

Probiotics are living microorganisms that provide a health benefit to the host when given in sufficient concentrations (Hill et al., 2014). The majority of *Lactobacillus*, *Bifidobacterium*, *Saccharomyces*, and *Bacillus* are the most often utilized probiotics in supplements, while *Lactobacillus* is the most generally used. *Bifidobacterium* and *Lactobacillus acidophilus* are solely utilized in foods (Antunes et al., 2020).

Probiotics, in combination with a tailored diet, may help to balance the microbiota and boost the host's immune system function. Probiotics may be able to moderate amplified immunological responses, such as those caused by antibiotics and the cytokine storm caused by COVID-19. Probiotics may decrease local and systemic inflammatory reactions in COVID-19 by minimizing the cytokine storm on the axis of the stomach, lungs, and brain (Ailioaie & Litscher, 2021). Modern therapeutic approaches have been substantially influenced

TABLE 2 Effects of probiotic and symbiotic supplementation on human health

Supplement	Bacterial strain	Subject	Outcomes	References
Probiotic	<i>Bifidobacterium longum</i>	Adults	High niche difference was observed and vacant niches were available when there were no competitors for externally added microbes	Maldonado-Gómez et al. (2016)
Probiotics	<i>Bifidobacterium infantis</i>	Infants	EVC001 had a clear advantage due to the presence of HMOs in breastmilk, and its administration along with breastfeeding encourages optimum conditions for a niche. EVC001 prevents the growth of gram-positive bacteria by lowering the pH, moreover HMOs encourage bacterial strain to persist in the infant gut	O'Brien et al. (2021) (Frese et al., 2017)
Probiotics	<i>L. rhamnosus</i> and <i>B. animalis</i>	Adults	During probiotic treatment, improvement in gut diversity was observed in 13% of patients with cystic fibrosis disease	Van Biervliet et al. (2018)
Probiotics and Insulin	<i>Lactobacillus rhamnosus</i>	Adults	Gut dysbiosis may reduce depression and chronic inflammation. The probiotic strain had shown antiinflammatory and antidepressant effects; particularly, cosupplementation with insulin in CAD patients had shown beneficial impacts	Bohlouli et al. (2021)
Probiotics	<i>Lactobacillus acidophilus</i> <i>Lactobacillus casei</i> <i>Bifidobacterium bifidum</i>	Women	Remarkable effects had been observed on glycemia, weight loss, and lipid profile of women with polycystic syndrome by probiotic strains supplementation	Ahmadi et al. (2017)
Probiotic	<i>L. acidophilus</i> <i>L. reuteri</i> <i>L. fermentum</i> <i>B. bifidum</i>	Women	Probiotic supplementation significantly improves weight loss, cholesterol, insulin, and hormonal balance in PCOs patients	Tabrizi et al. (2022)
Probiotic	<i>Lactobacillus pentosus</i>	Female BALB/c mice	Remarkable NK-cell activity was observed along with higher expression of IL-12 and IFN in lungs, and the endurance rate was high as compared to viral load in lungs with respect to influenza virus infection	Izumo et al. (2010)
Probiotic	<i>Lactobacillus rhamnosus</i>	Female BALB/c mice	Infected mice's clinical symptoms were relieved in response to influenza virus, and the viral burden in their lungs is dramatically reduced	Kawase et al. (2010)
Probiotic	<i>Lactobacillus rhamnosus</i>	6269 participants	According to meta-analysis of 23 studies, probiotics decrease the severity of pulmonary tract infections and improve mortality rate	Wang et al. (2016)
Probiotic	<i>Lactobacillus casei</i>	Adult/Children	Probiotics were found to be useful in the treatment of antibiotic-associated diarrhea	Dietrich et al. (2014)
Probiotic	<i>Lactobacillus rhamnosus</i>	Participants	A meta-analysis of 52 trials had shown that probiotics are effective against multidiseases like acute respiratory tract infections, antibiotic-associated diarrhea, acute infectious diarrhea, baby colic, and necrotizing enterocolitis	Liu et al. (2018)
Probiotic	<i>Lactobacillus gasseri</i>	Female BALB/c mice	Decreased expressions for proinflammatory cytokines, reduced weight loss, and viral load in the lungs of infected mice were observed	Eguchi et al. (2019)
Probiotic	<i>Lactobacillus rhamnosus</i>	Participants	A meta-analysis of 17 trials had proved that antibiotic use and the incidence of common acute illnesses were both dramatically decreased when probiotic strains were used	King et al. (2019)
Synbiotic	Combined strains	5 trials, 1049 Participants	A meta-analysis implies that synbiotic intervention can help individuals with metabolic syndrome reduce their cardiometabolic risk factors. This study had shown that how a synbiotic intervention affected people's lipid profiles, insulin resistance, blood pressure, anthropometric measurements, and inflammatory markers	Piper et al. (2022)

TABLE 2 (Continued)

Supplement	Bacterial strain	Subject	Outcomes	References
Synbiotic Probiotic Prebiotic	Strains of genera used: <i>Lactobacillus</i> <i>Bifidobacterium</i> <i>Propionibacterium</i> <i>Streptococcus</i>	Pregnant and lactating women	The impact of symbiotic formulas was observed with the combination of probiotic strains and galacto-oligosaccharides, fructo-oligosaccharides, or bovine milk-derived oligosaccharides Beneficial effects were observed in vaginally delivered, cesarean-delivered, and breastfed newborns	Hertz et al. (2022)
Probiotic	<i>Streptococcus thermophiles</i> <i>Lactobacillus acidophilus</i> <i>Lactobacillus helveticus</i> <i>Lactocaseibacillus paracasei</i> <i>Lactiplantibacillus plantarum</i> <i>Levilactobacillus brevis</i> <i>Bifidobacterium lactis</i> <i>Bifidobacterium lactis</i> <i>Lactocaseibacillus rhamnosus</i>	Hospitalized Covid-19 patients	An intervention of 2.4 billion CFU bacteria was given orally to 70 hospitalized Covid-19 patients. The clinical circumstances of COVID-19 patients improved significantly after a probiotic intervention	d'Ettorre et al. (2020)
Probiotic	Multiple strains	Covid-19 cases	A retrospective study was conducted on 800 positive Covid-19 cases to check the remission of diarrhea in Covid-19 patients. In comparison to the placebo group, the duration of diarrhea in the probiotic group was much shorter. Individuals' gastrointestinal symptoms such as abdominal distension, nausea, vomiting, and others were reduced as a result of the various strains	Xavier-Santos et al. (2022)
Probiotic	<i>Ligilactobacillus salivarius</i>	Covid-19-positive cases	29 COVID-positive residents of a nursing home were intervened daily with probiotic strain per unit of 125g. The study proposed that certain immunological variables might be used as nasal or fecal biomarkers to assess the advantages of a probiotic strain supplementation in the diet of elderly patients infected with SARS-CoV-2	Mozota et al. (2021)

by the interaction between the GM and host immune system, and altering the gut microbiota (GM) to improve the acquired immunological response in aged people is drawing interest. Diversification of gut microbiota with probiotics, prebiotics, or postbiotics may be an effective method to improve the effects of immunization in immune-compromised elderly people to boost immunological responses, including long-term NK-cell activity and antibody titers, as well as restore the GM balance (Akatsu, 2021).

The gut microbiota also has a significant impact on immunity and inflammation. It also has an effect on pulmonary function via the gut-lung axis. The role of the host microbiome in infection and pathogenicity has recently been reported. Understanding the microbiomes of the stomach and lungs might lead to novel medicinal methods (Allali et al., 2021). Nutritional techniques to boost immunity against SARS-CoV-2 are being considered as most promising techniques by various studies. Prebiotics, on the other hand, may improve gut immunity by boosting particular gut microorganisms selectively. Probiotics, prebiotics, and fermented foods have been shown to improve gut and pulmonary health along with immunity at various levels of proof. Without a guarantee of efficacy against

COVID-19, the inclusion of such nutritional foods in the diet may reduce gut inflammation, reduce the intensity of infection, and boost mucosal immunity (Antunes et al., 2020).

COVID-19 has been demonstrated to have impacts on gut microbiota dysbiosis, and probiotics have shown a remarkable role in the treatment of pulmonary tract viral infections. These findings may boost the impact of probiotics as a treatment therapy for COVID-19 (Bottari et al., 2021). Oral probiotics, which are beneficial living bacteria that can give medicinal benefits to the host when provided in safe doses, may be worth examining as a potential preventative or therapeutic method to minimize coronavirus illness and the intensity of the symptoms (Baindara et al., 2021). The effects of probiotic and symbiotic supplementation on human health are presented in Table 2.

6 | CONCLUSION

The gut microbiota is widely known for providing a variety of health benefits, including disease protection. Although gut dysbiosis, which can be induced by a variety of factors, is linked to severe COVID-19,

it is essential to address the approaches to maintain the health of gut microbiota. Diet affects the makeup and functional capabilities of the gut microbiota, which in turn affects host biochemical processes, forming a system of mutual interaction and interdependence. Macronutrients, fiber, polyphenols, and prebiotics are all important factors in shaping the composition of gut microbiota. COVID-19 is less severe when the gut microbiota is healthy, diverse, and abundant, as well it also increases the efficacy of vaccinations against COVID-19.

ACKNOWLEDGMENT

The authors are thankful to Government College University for providing literature collection facilities.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

DATA AVAILABILITY STATEMENT

Even though adequate data have been given in the form of tables and figures, all authors declare that if more data are required, then the data will be provided on a request basis.

ETHICS STATEMENT

This article does not contain any studies with human participants or animals performed by any of the authors.

CONSENT TO PARTICIPATE

Corresponding and all the coauthors are willing to participate in this manuscript.

CONSENT FOR PUBLICATION

All authors are willing to the publication of this manuscript.

ORCID

Farhan Saeed  <https://orcid.org/0000-0001-5340-4015>

Yasir Abbas Shah  <https://orcid.org/0000-0003-4925-459X>

Huda Ateeq  <https://orcid.org/0000-0001-9612-1031>

Muhammad Nadeem Akhtar  <https://orcid.org/0000-0002-8610-2686>

[org/0000-0002-8610-2686](https://orcid.org/0000-0002-8610-2686)

Tabassum Tufail  <https://orcid.org/0000-0002-7632-5261>

REFERENCES

- Acosta-Elias, J., & Espinosa-Tanguma, R. (2020). The folate concentration and/or folic acid metabolites in plasma as factor for COVID-19 infection. *Frontiers in Pharmacology*, *11*, 1062.
- Ahmadi, S., Jamilian, M., Karamali, M., Tajabadi-Ebrahimi, M., Jafari, P., Taghizadeh, M., Memarzadeh, M. R., & Asemi, Z. (2017). Probiotic supplementation and the effects on weight loss, glycaemia and lipid profiles in women with polycystic ovary syndrome: A randomized, double-blind, placebo-controlled trial. *Human Fertility*, *20*(4), 254–261.
- Ailioaie, L. M., & Litscher, G. (2021). Probiotics, photobiomodulation, and disease management: Controversies and challenges. *International Journal of Molecular Sciences*, *22*(9), 4942.
- Akatsu, H. (2021). Exploring the effect of probiotics, prebiotics, and postbiotics in strengthening immune activity in the elderly. *Vaccine*, *9*(2), 136.
- Alameddine, J., Godefroy, E., Papargyris, L., Sarrabayrouse, G., Tabiasco, J., Bridonneau, C., Yazdanbakhsh, K., Sokol, H., Altare, F., & Jotereau, F. (2019). *Faecalibacterium prausnitzii* skews human DC to prime IL10-producing T cells through TLR2/6/JNK signaling and IL-10, IL-27, CD39, and IDO-1 induction. *Frontiers in Immunology*, *10*, 143.
- Alberca, R. W., Oliveira, L. D. M., Branco, A. C. C. C., Pereira, N. Z., & Sato, M. N. (2021). Obesity as a risk factor for COVID-19: An overview. *Critical Reviews in Food Science and Nutrition*, *61*(13), 2262–2276.
- Allali, I., Bakri, Y., Amzazi, S., & Ghazal, H. (2021). Gut-lung axis in COVID-19. *Interdisciplinary Perspectives on Infectious Diseases*, *2021*, 1–6.
- Amaral, J. F. D., Foschetti, D. A., Assis, F. A. D., Menezes, J. S., Vaz, N. M., & Faria, A. M. C. D. (2006). Immunoglobulin production is impaired in protein-deprived mice and can be restored by dietary protein supplementation. *Brazilian Journal of Medical and Biological Research*, *39*(12), 1581–1586.
- Anand, S., & Mande, S. S. (2018). Diet, microbiota and gut-lung connection. *Frontiers in Microbiology*, *9*, 2147.
- Antunes, A. E., Vinderola, G., Xavier-Santos, D., & Sivieri, K. (2020). Potential contribution of beneficial microbes to face the COVID-19 pandemic. *Food Research International*, *136*, 109577.
- Bae, S., Kim, S. R., Kim, M. N., Shim, W. J., & Park, S. M. (2021). Impact of cardiovascular disease and risk factors on fatal outcomes in patients with COVID-19 according to age: A systematic review and meta-analysis. *Heart*, *107*(5), 373–380.
- Baindara, P., Chakraborty, R., Holliday, Z. M., Mandal, S. M., & Schrum, A. G. (2021). Oral probiotics in coronavirus disease 2019: Connecting the gut–lung axis to viral pathogenesis, inflammation, secondary infection and clinical trials. *New Microbes and New Infections*, *40*, 100837.
- Bajinka, O., Simbilyabo, L., Tan, Y., Jabang, J., & Saleem, S. A. (2022). Lung-brain axis. *Critical Reviews in Microbiology*, *48*(3), 257–269.
- Barnett, J. B., Hamer, D. H., & Meydani, S. N. (2010). Low zinc status: A new risk factor for pneumonia in the elderly? *Nutrition Reviews*, *68*(1), 30–37.
- Bingula, R., Filaire, M., Radosevic-Robin, N., Bey, M., Berthon, J. Y., Bernalier-Donadille, A., Vasson, M. P., & Filaire, E. (2017). Desired turbulence? Gut-lung axis immunity and lung cancer. *Journal of Oncology*, *2017*, 5035371.
- Bohlouli, J., Namjoo, I., Borzoo-Isfahani, M., Kermani, M. A. H., Zehi, Z. B., & Moravejolahkami, A. R. (2021). Effect of probiotics on oxidative stress and inflammatory status in diabetic nephropathy: A systematic review and meta-analysis of clinical trials. *Heliyon*, *7*(1), e05925.
- Bottari, B., Castellone, V., & Neviani, E. (2021). Probiotics and COVID-19. *International Journal of Food Sciences and Nutrition*, *72*(3), 293–299.
- Bouhnik, Y., Achour, L., Paineau, D., Riottot, M., Attar, A., & Bornet, F. (2007). Four-week short chain fructo-oligosaccharides ingestion leads to increasing fecal bifidobacteria and cholesterol excretion in healthy elderly volunteers. *Nutrition Journal*, *6*(1), 1–7.
- Callaway, E. (2021). COVID vaccine boosters: The most important questions. *Nature*, *596*(7871), 178–180.
- Carr, A. C., & Maggini, S. (2017). Vitamin C and immune function. *Nutrients*, *9*(11), 1211.
- Carr, N. B., & Melcher, C. P. (2017). *Wyoming Basin rapid ecoregional assessment*. US Department of the Interior, US Geological Survey.
- Cheng, R. Z. (2020). Can early and high intravenous dose of vitamin C prevent and treat coronavirus disease 2019 (COVID-19)? *Medicine in Drug Discovery*, *5*, 100028.
- Cheung, K. S., Hung, I. F., Chan, P. P., Lung, K. C., Tso, E., Liu, R., Ng, Y. Y., Chu, M. Y., TWH, C., Tam, A. R., CCY, Y., Leung, K. H., Fung, A. Y., Zhang, R. R., Lin, Y., Cheng, H. M., AJX, Z., To KKW, Chan, K. H., ... Leung, W. K. (2020). Gastrointestinal manifestations of SARS-CoV-2 infection and virus load in fecal samples from a Hong Kong

- cohort: Systematic review and meta-analysis. *Gastroenterology*, 159(1), 81–95.
- Childs, C. E., Calder, P. C., & Miles, E. A. (2020). Diet and immune function. *Nutrients*, 2019(11), 1933.
- Chowdhury, M. A., Hossain, N., Kashem, M. A., Shahid, M. A., & Alam, A. (2020). Immune response in COVID-19: A review. *Journal of Infection and Public Health*, 13(11), 1619–1629.
- Chungchunlam, S. M., Henare, S. J., Ganesh, S., & Moughan, P. J. (2017). Effects of whey protein and its two major protein components on satiety and food intake in normal-weight women. *Physiology & Behavior*, 175, 113–118.
- De Santis, S., Cavalcanti, E., Mastronardi, M., Jirillo, E., & Chieppa, M. (2015). Nutritional keys for intestinal barrier modulation. *Frontiers in Immunology*, 6, 612.
- d'Ettorre, G., Ceccarelli, G., Marazzato, M., Campagna, G., Pinacchio, C., Alessandri, F., Ruberto, F., Rossi, G., Celani, L., Scagnolari, C., Mastropietro, C., Trinchieri, V., Recchia, G. E., Mauro, V., Antonelli, G., Pugliese, F., & Mastroianni, C. M. (2020). Challenges in the management of SARS-CoV2 infection: The role of oral bacteriotherapy as complementary therapeutic strategy to avoid the progression of COVID-19. *Frontiers in Medicine*, 7, 389.
- Dhar, D., & Mohanty, A. (2020). Gut microbiota and Covid-19-possible link and implications. *Virus Research*, 285, 198018.
- Dietrich, C. G., Kottmann, T., & Alavi, M. (2014). Commercially available probiotic drinks containing lactobacillus casei DN-114001 reduce antibiotic-associated diarrhea. *World Journal of Gastroenterology: WJG*, 20(42), 15837–15844.
- Dominika, Š., Arjan, N., Karyn, R. P., & Henryk, K. (2011). The study on the impact of glycosylated pea proteins on human intestinal bacteria. *International Journal of Food Microbiology*, 145(1), 267–272.
- Dubourg, G., Edouard, S., & Raoult, D. (2019). Relationship between nasopharyngeal microbiota and patient's susceptibility to viral infection. *Expert Review of Anti-Infective Therapy*, 17(6), 437–447.
- Dumas, A., Bernard, L., Poquet, Y., Lugo-Villarino, G., & Neyrolles, O. (2018). The role of the lung microbiota and the gut–lung axis in respiratory infectious diseases. *Cellular Microbiology*, 20(12), e12966.
- Earley, Z. M., Akhtar, S., Green, S. J., Naqib, A., Khan, O., Cannon, A. R., Hammer, A. M., Morris, N. L., Li, X., Eberhardt, J. M., Gamelli, R. L., Kennedy, R. H., & Choudhry, M. A. (2015). Burn injury alters the intestinal microbiome and increases gut permeability and bacterial translocation. *PLoS One*, 10(7), e0129996.
- Eguchi, K., Fujitani, N., Nakagawa, H., & Miyazaki, T. (2019). Prevention of respiratory syncytial virus infection with probiotic lactic acid bacterium *Lactobacillus gasseri* SBT2055. *Scientific Reports*, 9, 1–11.
- Enaud, R., Prevel, R., Ciarlo, E., Beauflis, F., Wieërs, G., Guery, B., & Delhaes, L. (2020). The gut-lung axis in health and respiratory diseases: A place for inter-organ and inter-kingdom crosstalks. *Frontiers in Cellular and Infection Microbiology*, 10, 9.
- Fanos, V., Pintus, M. C., Pintus, R., & Marcialis, M. A. (2020). Lung microbiota in the acute respiratory disease: From coronavirus to metabolomics. *Journal of Pediatric and Neonatal Individualized Medicine (JPNIM)*, 9(1), e090139.
- Frese, S. A., Hutton, A. A., Contreras, L. N., Shaw, C. A., Palumbo, M. C., Casaburi, G., Xu, G., JCC, D., Lebrilla, C. B., Henrick, B. M., Freeman, S. L., Barile, D., German, J. B., Mills, D. A., Smilowitz, J. T., & Underwood, M. A. (2017). Persistence of supplemented *Bifidobacterium longum* subsp. infantis EVC001 in breastfed infants. *MSphere*, 2(6), e00501-17.
- Galdeano, C. M., Cazorla, S. I., Dumit, J. M. L., Vélez, E., & Perdígón, G. (2019). Beneficial effects of probiotic consumption on the immune system. *Annals of Nutrition and Metabolism*, 74(2), 115–124.
- Galland, L. (2010). Diet and inflammation. *Nutrition in Clinical Practice*, 25(6), 634–640.
- Gill, S. R., Pop, M., Deboy, R. T., Eckburg, P. B., Turnbaugh, P. J., Samuel, B. S., Gordon, J. I., Relman, D. A., Fraser-Liggett, C. M., & Nelson, K. E. (2006). Metagenomic analysis of the human distal gut microbiome. *Science*, 312(5778), 1355–1359.
- Grant, W. B., Lahore, H., McDonnell, S. L., Baggerly, C. A., French, C. B., Aliano, J. L., & Bhattoa, H. P. (2020). Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. *Nutrients*, 12(4), 988.
- Groves, H. T., Higham, S. L., Moffatt, M. F., Cox, M. J., & Tregoning, J. S. (2020). Respiratory viral infection alters the gut microbiota by inducing inappetence. *MBio*, 11(1), e03236–e03219.
- Guaraldi, F., & Salvatori, G. (2012). Effect of breast and formula feeding on gut microbiota shaping in newborns. *Frontiers in Cellular and Infection Microbiology*, 2, 94.
- Hagan, T., Cortese, M., Roupheal, N., Boudreau, C., Linde, C., Maddur, M. S., Das, J., Wang, H., Guthmiller, J., Zheng, N. Y., Huang, M., Uphadhyay, A. A., Gardinassi, L., Petittedmange, C., MP, M. C., Johnson, S. J., Gill, K., Cervasi, B., Zou, J., ... Pulendran, B. (2019). Antibiotics-driven gut microbiome perturbation alters immunity to vaccines in humans. *Cell*, 178(6), 1313–1328.
- Hahn, O., Grönke, S., Stubbs, T. M., Ficz, G., Hendrich, O., Krueger, F., Andrews, S., Zhang, Q., Wakelam, M. J., Beyer, A., Reik, W., & Partridge, L. (2017). Dietary restriction protects from age-associated DNA methylation and induces epigenetic reprogramming of lipid metabolism. *Genome Biology*, 18(1), 1–18.
- Hall, A. B., Tolonen, A. C., & Xavier, R. J. (2017). Human genetic variation and the gut microbiome in disease. *Nature Reviews Genetics*, 18(11), 690–699.
- Hall, J. A., Grainger, J. R., Spencer, S. P., & Belkaid, Y. (2011). The role of retinoic acid in tolerance and immunity. *Immunity*, 35(1), 13–22.
- Hemilä, H. (2017). Vitamin C and infections. *Nutrients*, 9(4), 339.
- Hertz, S., Durack, J., Kirk, K. F., Nielsen, H. L., Lin, D. L., Fadros, D., Lynch, K., Piceno, Y., Thorlacius-Ussing, O., Nielsen, H., & Lynch, S. V. (2022). Microscopic colitis patients possess a perturbed and inflammatory gut microbiota. *Digestive Diseases and Sciences*, 67(6), 2433–2443.
- Hill, C., Guarner, F., Reid, G., Gibson, G. R., Merenstein, D. J., Pot, B., Morelli, L., Canani, R. B., Flint, H. J., Salminen, S., Calder, P. C., & Sanders, M. E. (2014). Expert consensus document: The international scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nature Reviews Gastroenterology & Hepatology*, 11(8), 506–514.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., ... Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*, 395(10223), 497–506.
- Huang, Z., Liu, Y., Qi, G., Brand, D., & Zheng, S. G. (2018). Role of vitamin A in the immune system. *Journal of Clinical Medicine*, 7(9), 258.
- Human Microbiome Project Consortium. (2012). A framework for human microbiome research. *Nature*, 486(7402), 215–221.
- Iddir, M., Brito, A., Dinger, G., Fernandez Del Campo, S. S., Samouda, H., La Frano, M. R., & Bohn, T. (2020). Strengthening the immune system and reducing inflammation and oxidative stress through diet and nutrition: Considerations during the COVID-19 crisis. *Nutrients*, 12(6), 1562.
- Innes, J. K., & Calder, P. C. (2018). Omega-6 fatty acids and inflammation. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 132, 41–48.
- Izumo, T., Maekawa, T., Ida, M., Noguchi, A., Kitagawa, Y., Shibata, H., Yasui, H., & Kiso, Y. (2010). Effect of intranasal administration of *Lactobacillus pentosus* S-PT84 on influenza virus infection in mice. *International Immunopharmacology*, 10(9), 1101–1106.
- Kaminogawa, S., & Nanno, M. (2004). Modulation of immune functions by foods. *Evidence-based Complementary and Alternative Medicine*, 1(3), 241–250.
- Kanauchi, O., Andoh, A., AbuBakar, S., & Yamamoto, N. (2018). Probiotics and paraprobiotics in viral infection: Clinical application and effects

- on the innate and acquired immune systems. *Current Pharmaceutical Design*, 24(6), 710–717.
- Kawase, M., He, F., Kubota, A., Harata, G., & Hiramatsu, M. (2010). Oral administration of lactobacilli from human intestinal tract protects mice against influenza virus infection. *Letters in Applied Microbiology*, 51(1), 6–10.
- Keely, S., Talley, N. J., & Hansbro, P. M. (2012). Pulmonary-intestinal cross-talk in mucosal inflammatory disease. *Mucosal Immunology*, 5(1), 7–18.
- Kilgore, P. (2020). *The COVID-19 survival guide: Information and practical steps you can use to protect yourself against the coronavirus*. Quantitative Health, LLC.
- King, S., Tancredi, D., Lenoir-Wijnkoop, I., Gould, K., Vann, H., Connors, G., Sanders, M. E., Linder, J. A., Shane, A. L., & Merenstein, D. (2019). Does probiotic consumption reduce antibiotic utilization for common acute infections? A systematic review and meta-analysis. *European Journal of Public Health*, 29(3), 494–499.
- Lake, M. A. (2020). What we know so far: COVID-19 current clinical knowledge and research. *Clinical Medicine*, 20(2), 124–127.
- Lau, H. C. H., & Yu, J. (2022). Targeting the gut microbiota in coronavirus disease 2019: Hype or Hope? *Gastroenterology*, 162(1), 9–16.
- Lauridsen, C. (2019). From oxidative stress to inflammation: Redox balance and immune system. *Poultry Science*, 98(10), 4240–4246.
- Le Chatelier, E., Nielsen, T., Qin, J., Prifti, E., Hildebrand, F., Falony, G., Almeida, M., Arumugam, M., Batto, J. M., Kennedy, S., Leonard, P., Li, J., Burgdorf, K., Grarup, N., Jørgensen, T., Brandslund, I., Nielsen, H. B., Juncker, A. S., Bertalan, M., ... Pedersen, O. (2013). Richness of human gut microbiome correlates with metabolic markers. *Nature*, 500(7464), 541–546.
- Li, X., He, J., Yu, M., & Sun, J. (2020). The efficacy of vitamin D therapy for patients with COPD: A meta-analysis of randomized controlled trials. *Annals of Palliative Medicine*, 9, 286–297.
- Lindgren, E., Harris, F., Dangour, A. D., Gasparatos, A., Hiramatsu, M., Javadi, F., Loken, B., Murakami, T., Scheelbeek, P., & Haines, A. (2018). Sustainable food systems—A health perspective. *Sustainability Science*, 13(6), 1505–1517.
- Liu, Y., Tran, D. Q., & Rhoads, J. M. (2018). Probiotics in disease prevention and treatment. *The Journal of Clinical Pharmacology*, 58, S164–S179.
- Maldonado-Gómez, M. X., Martínez, I., Bottacini, F., O'Callaghan, A., Ventura, M., van Sinderen, D., Hillmann, B., Vangay, P., Knights, D., Hutkins, R. W., & Walter, J. (2016). Stable engraftment of *Bifidobacterium longum* AH1206 in the human gut depends on individualized features of the resident microbiome. *Cell Host & Microbe*, 20(4), 515–526.
- Moens, E., & Veldhoen, M. (2012). Epithelial barrier biology: Good fences make good neighbours. *Immunology*, 135(1), 1–8.
- Mosca, A., Leclerc, M., & Hugot, J. P. (2016). Gut microbiota diversity and human diseases: Should we reintroduce key predators in our ecosystem? *Frontiers in Microbiology*, 7, 455.
- Mosekilde, L. (2005). Vitamin D and the elderly. *Clinical Endocrinology*, 62(3), 265–281.
- Mozota, M., Castro, I., Gómez-Torres, N., Arroyo, R., Laila, Y., Somada, M., Alba, C., & Rodríguez, J. M. (2021). Administration of *Ligilactobacillus salivarius* MP101 in an elderly nursing home during the COVID-19 pandemic: Immunological and nutritional impact. *Food*, 10(9), 2149.
- Nagpal, R., Mainali, R., Ahmadi, S., Wang, S., Singh, R., Kavanagh, K., Kitzman, D. W., Kushugulova, A., Marotta, F., & Yadav, H. (2018). Gut microbiome and aging: Physiological and mechanistic insights. *Nutrition and Healthy Aging*, 4(4), 267–285.
- Negi, S., Pahari, S., Bashir, H., & Agrewala, J. N. (2019). Gut microbiota regulates mIncle mediated activation of I ung dendritic cells to protect against *Mycobacterium tuberculosis*. *Frontiers in Immunology*, 10, 1142.
- O'Brien, C. E., Meier, A. K., Cernioglio, K., Mitchell, R. D., Casaburi, G., Frese, S. A., Henrick, B. M., Underwood, M. A., & Smilowitz, J. T. (2022). Early probiotic supplementation with *B. infantis* in breastfed infants leads to persistent colonization at 1 year. *Pediatric Research*, 91, 627–636.
- O'Keefe, J. H., Gheewala, N. M., & O'Keefe, J. O. (2008). Dietary strategies for improving post-prandial glucose, lipids, inflammation, and cardiovascular health. *Journal of the American College of Cardiology*, 51(3), 249–255.
- Parada Venegas, D., De la Fuente, M. K., Landskron, G., González, M. J., Quera, R., Dijkstra, G., HJM, H., Faber, K. N., & Hermoso, M. A. (2019). Short chain fatty acids (SCFAs)-mediated gut epithelial and immune regulation and its relevance for inflammatory bowel diseases. *Frontiers in Immunology*, 277.
- Park, Y., Subar, A. F., Hollenbeck, A., & Schatzkin, A. (2011). Dietary fiber intake and mortality in the NIH-AARP diet and health study. *Archives of Internal Medicine*, 171(12), 1061–1068.
- Parker, E. P., Kampmann, B., Kang, G., & Grassly, N. C. (2014). Influence of enteric infections on response to oral poliovirus vaccine: A systematic review and meta-analysis. *The Journal of Infectious Diseases*, 210(6), 853–864.
- Piper, S., Jiwani, S. I., Gyasi-Antwi, P., & Adams, G. G. (2021). The benefits and risks of probiotic, prebiotic and symbiotic interventions in the care of patients with diabetes mellitus. *International Journal of Diabetes and Endocrinology*, 110, 10.
- Pormohammad, A., Monych, N. K., & Turner, R. J. (2021). Zinc and SARS-CoV-2: A molecular modeling study of Zn interactions with RNA-dependent RNA-polymerase and 3C-like proteinase enzymes. *International Journal of Molecular Medicine*, 47(1), 326–334.
- Pulendran, B. (2014). Systems vaccinology: Probing humanity's diverse immune systems with vaccines. *Proceedings of the National Academy of Sciences of the United States of America*, 111(34), 12300–12306.
- Radzikowska, U., Rinaldi, A. O., Çelebi Sözen, Z., Karaguzel, D., Wojcik, M., Cypryk, K., Akdis, M., Akdis, C. A., & Sokolowska, M. (2019). The influence of dietary fatty acids on immune responses. *Nutrients*, 11(12), 2990.
- Raharusun, P., Priambada, S., Budiarti, C., Agung, E., & Budi, C. (2020). Patterns of COVID-19 mortality and vitamin D: An Indonesian study. *SSRN Electronic Journal*, 7, 1–12.
- Read, S. A., Obeid, S., Ahlenstiel, C., & Ahlenstiel, G. (2019). The role of zinc in antiviral immunity. *Advances in Nutrition*, 10(4), 696–710.
- Ren, M., Zhang, S. H., Zeng, X. F., Liu, H., & Qiao, S. Y. (2015). Branched-chain amino acids are beneficial to maintain growth performance and intestinal immune-related function in weaned piglets fed protein restricted diet. *Asian-Australasian Journal of Animal Sciences*, 28(12), 1742–1750.
- Robichaud, S., Shahbazi, R., & Matar, C. (2021). Role of probiotics in prevention of COVID-19 through modulation of gut-lung axis. In *COVID-19 and Nutraceuticals, A Guidebook: COVID-19 and Nutraceuticals* (p. 33).
- Rodríguez, L., Cervantes, E., & Ortiz, R. (2011). Malnutrition and gastrointestinal and respiratory infections in children: A public health problem. *International Journal of Environmental Research and Public Health*, 8(4), 1174–1205.
- Ross, A. C. (2010). Diet in vitamin A research. In J. M. Walker (Ed.), *Retinoids* (pp. 295–313). Humana Press.
- Saeed, F., Nadeem, M., Ahmed, R. S., Tahir Nadeem, M., Arshad, M. S., & Ullah, A. (2016). Studying the impact of nutritional immunology underlying the modulation of immune responses by nutritional compounds—A review. *Food and Agricultural Immunology*, 27(2), 205–229.
- Schoeler, M., & Caesar, R. (2019). Dietary lipids, gut microbiota and lipid metabolism. *Reviews in Endocrine and Metabolic Disorders*, 20(4), 461–472.
- Serhan, C. N., & Levy, B. D. (2018). Resolvins in inflammation: Emergence of the pro-resolving superfamily of mediators. *Journal of Clinical Investigation*, 128, 2657–2669.
- Shrestha, S., Kim, S. Y., Yun, Y. J., Kim, J. K., Lee, J. M., Shin, M., Song, D. K., & Hong, C. W. (2017). Retinoic acid induces hypersegmentation

- and enhances cytotoxicity of neutrophils against cancer cells. *Immunology Letters*, 182, 24–29.
- Siddiqi, H. K., & Mehra, M. R. (2020). COVID-19 illness in native and immunosuppressed states: A clinical– Therapeutic staging proposal. *The Journal of Heart and Lung Transplantation*, 39(5), 405–407.
- Singh, B., Read, S., Asseman, C., Malmström, V., Mottet, C., Stephens, L. A., Stepankova, R., Tlaskalova, H., & Powrie, F. (2001). Control of intestinal inflammation by regulatory T cells. *Immunological Reviews*, 182(1), 190–200.
- Stefanidou, M., Maravelias, C., Dona, A., & Spiliopoulou, C. (2006). Zinc: a multipurpose trace element. *Archives of Toxicology*, 80(1), 1–9.
- Tabrizi, R., Ostadmohammadi, V., Akbari, M., Lankarani, K. B., Vakili, S., Peymani, P., Karamali, M., Kolahdooz, F., & Asemi, Z. (2022). The effects of probiotic supplementation on clinical symptom, weight loss, glycemic control, lipid and hormonal profiles, biomarkers of inflammation, and oxidative stress in women with polycystic ovary syndrome: A systematic review and meta-analysis of randomized controlled trials. *Probiotics and Antimicrobial Proteins*, 14(1), 1–14.
- Taylor, A. K., Cao, W., Vora, K. P., Cruz, J. D. L., Shieh, W. J., Zaki, S. R., Katz, J. M., Sambhara, S., & Gangappa, S. (2013). Protein energy malnutrition decreases immunity and increases susceptibility to influenza infection in mice. *The Journal of Infectious Diseases*, 207(3), 501–510.
- Timoneda, J., Rodríguez-Fernández, L., Zaragoza, R., Marín, M. P., Cabezuolo, M. T., Torres, L., Viña, J. R., & Barber, T. (2018). Vitamin A deficiency and the lung. *Nutrients*, 10(9), 1132.
- Trompette, A., Gollwitzer, E. S., Pattaroni, C., Lopez-Mejia, I. C., Riva, E., Pernot, J., Ubags, N., Fajas, L., Nicod, L. P., & Marsland, B. J. (2018). Dietary fiber confers protection against flu by shaping Ly6c⁺ patrolling monocyte hematopoiesis and CD8⁺ T cell metabolism. *Immunity*, 48(5), 992–1005.
- Truwit, J. D., Hite, R. D., Morris, P. E., DeWilde, C., Priday, A., Fisher, B., Thacker, L. R., 2nd, Natarajan, R., Brophy, D. F., Sculthorpe, R., Nanchal, R., Syed, A., Sturgill, J., Martin, G. S., Sevransky, J., Kashiouris, M., Hamman, S., Egan, K. F., Hastings, A., ... Halquist, M. (2019). Effect of vitamin C infusion on organ failure and biomarkers of inflammation and vascular injury in patients with sepsis and severe acute respiratory failure: The CITRIS-ALI randomized clinical trial. *JAMA*, 322(13), 1261–1270.
- Valdez, Y., Brown, E. M., & Finlay, B. B. (2014). Influence of the microbiota on vaccine effectiveness. *Trends in Immunology*, 35(11), 526–537.
- Van Biervliet, S., Hauser, B., Verhulst, S., Stepman, H., Delanghe, J., Warzee, J. P., Vandewiele, T., & Wilschanski, M. (2018). Probiotics in cystic fibrosis patients: A double blind crossover placebo controlled study: Pilot study from the ESPGHAN working group on pancreas/CF. *Clinical Nutrition ESPEN*, 27, 59–65.
- Villanueva-Millán, M. J., Pérez-Matute, P., & Oteo, J. A. (2015). Gut microbiota: A key player in health and disease. A review focused on obesity. *Journal of physiology and biochemistry*, 71(3), 509–525.
- Vlasova, A. N., Takanashi, S., Miyazaki, A., Rajashekar, G., & Saif, L. J. (2019). How the gut microbiome regulates host immune responses to viral vaccines. *Current Opinion in Virology*, 37, 16–25.
- Voigt, R. M., Forsyth, C. B., Green, S. J., Mutlu, E., Engen, P., Vitaterna, M. H., Turek, F. W., & Keshavarzian, A. (2014). Circadian disorganization alters intestinal microbiota. *PLoS One*, 9(5), e97500.
- Von Essen, M. R., Kongsbak, M., Schjerling, P., Olgaard, K., Ødum, N., & Geisler, C. (2010). Vitamin D controls T cell antigen receptor signaling and activation of human T cells. *Nature Immunology*, 11(4), 344–349.
- Wang, Y., Li, X., Ge, T., Xiao, Y., Liao, Y., Cui, Y., Zhang, Y., Ho, W., Yu, G., & Zhang, T. (2016). Probiotics for prevention and treatment of respiratory tract infections in children: A systematic review and meta-analysis of randomized controlled trials. *Medicine*, 95(31), e4509.
- Wannamethee, S. G., Lowe, G. D., Rumley, A., Bruckdorfer, K. R., & Whincup, P. H. (2006). Associations of vitamin C status, fruit and vegetable intakes, and markers of inflammation and hemostasis. *The American Journal of Clinical Nutrition*, 83(3), 567–574.
- Watchorn, J., Clasky, A. J., Prakash, G., Johnston, I. A., Chen, P. Z., & Gu, F. X. (2022). Untangling mucosal drug delivery: Engineering, designing, and testing nanoparticles to overcome the mucus barrier. *ACS Biomaterials Science & Engineering*, 8(4), 1396–1426.
- Wessells, K. R., & Brown, K. H. (2012). Estimating the global prevalence of zinc deficiency: Results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One*, 7(11), e50568.
- Wessels, I., Maywald, M., & Rink, L. (2017). Zinc as a gatekeeper of immune function. *Nutrients*, 9(12), 1286.
- Wintergerst, E. S., Maggini, S., & Hornig, D. H. (2006). Immune-enhancing role of vitamin C and zinc and effect on clinical conditions. *Annals of Nutrition and Metabolism*, 50(2), 85–94.
- Wintergerst, E. S., Maggini, S., & Hornig, D. H. (2007). Contribution of selected vitamins and trace elements to immune function. *Annals of Nutrition and Metabolism*, 51(4), 301–323.
- Wirth, J. P., Petry, N., Tanumihardjo, S. A., Rogers, L. M., McLean, E., Greig, A., Garrett, G. S., Klemm, R. D., & Rohner, F. (2017). Vitamin A supplementation programs and country-level evidence of vitamin A deficiency. *Nutrients*, 9(3), 190.
- Xavier-Santos, D., Padilha, M., Fabiano, G. A., Vinderola, G., Cruz, A. G., Sivieri, K., & Antunes, A. E. C. (2022). Corrigendum to 'Evidences and perspectives of the use of probiotics, prebiotics, synbiotics, and postbiotics as adjuvants for prevention and treatment of COVID-19: A bibliometric analysis and integrative review' [Trends in Food Science & Technology 120 (2022) 174–192]. *Trends in Food Science & Technology*, 121, 156.
- Yan, J., & Horng, T. (2020). Lipid metabolism in regulation of macrophage functions. *Trends in Cell Biology*, 30(12), 979–989.
- Yang, Y., Shen, C., Li, J., Yuan, J., Yang, M., Wang, F., Li, G., Li, Y., Xing, L., Peng, L., & Liu, Y. (2020). Exuberant elevation of IP-10, MCP-3 and IL-1ra during SARS-CoV-2 infection is associated with disease severity and fatal outcome. *MedRxiv*. <https://doi.org/10.1016/j.jaci.2020.04.027>
- Yeoh, Y. K., Zuo, T., Lui, G. C. Y., Zhang, F., Liu, Q., Li, A. Y., Chung, A. C., Cheung, C. P., Tso, E. Y., Fung, K. S., Chan, V., Ling, L., Joynt, G., Hui, D. S., Chow, K. M., SSS, N., Li, T. C., Ng, R. W., Yip, T. C., ... Ng, S. C. (2021). Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19. *Gut*, 70(4), 698–706.
- Zhang, X., Liu, Y., Gu, Y., Wang, S., & Chen, H. (2020). Red for “stop”: “traffic-light” nutrition labels decrease unhealthy food choices by increasing activity and connectivity in the frontal lobe. *Nutrients*, 12(1), 128.
- Zhang, M., Zhu, J., Zhang, X., Zhao, D. G., Ma, Y. Y., Li, D., Ho, C. T., & Huang, Q. (2020). Aged citrus peel (chenpi) extract causes dynamic alteration of colonic microbiota in high-fat diet induced obese mice. *Food & Function*, 11(3), 2667–2678.
- Zuo, T., Zhang, F., Lui, G. C., Yeoh, Y. K., Li, A. Y., Zhan, H., Wan, Y., ACK, C., Cheung, C. P., Chen, N., CKC, L., Chen, Z., EYK, T., KSC, F., Chan, V., Ling, L., Joynt, G., DSC, H., FKL, C., ... Ng, S. C. (2020). Alterations in gut microbiota of patients with COVID-19 during time of hospitalization. *Gastroenterology*, 159(3), 944–955.

How to cite this article: Akram, N., Saeed, F., Afzaal, M., Shah, Y. A., Qamar, A., Faisal, Z., Ghani, S., Ateeq, H., Akhtar, M. N., Tufail, T., Hussain, M., Asghar, A., Rasheed, A., & Jbawi, E. A. (2023). Gut microbiota and synbiotic foods: Unveiling the relationship in COVID-19 perspective. *Food Science & Nutrition*, 11, 1166–1177. <https://doi.org/10.1002/fsn3.3162>