



## Correction: Species-differences in the in vitro biotransformation of trifluoroethene (HFO-1123)

R. Dekant<sup>1</sup> · R. Bertermann<sup>2</sup> · J. Serban<sup>1</sup> · S. Sharma<sup>1</sup> · M. Shinohara<sup>3</sup> · Y. Morizawa<sup>3</sup> · H. Okamoto<sup>3</sup> · W. Brock<sup>4</sup> · W. Dekant<sup>1</sup> · A. Mally<sup>1</sup> 

Published online: 6 December 2023  
© The Author(s) 2023

**Correction: Archives of Toxicology (2023) 97:3095–3111**  
<https://doi.org/10.1007/s00204-023-03603-3>

The original version of this article unfortunately contained a mistake. The presentation of Fig. 5 was incorrect. The corrected Fig. 5 is given below.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing,

adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright

---

The original article can be found online at <https://doi.org/10.1007/s00204-023-03603-3>.

---


✉ A. Mally  
mally@toxi.uni-wuerzburg.de

<sup>1</sup> Department of Toxicology, University of Würzburg, Versbacher Strasse 9, 97078 Würzburg, Germany

<sup>2</sup> Department of Inorganic Chemistry, University of Würzburg, Am Hubland, 97074 Würzburg, Germany

<sup>3</sup> Chemicals Company, AGC Inc, CSR Office, 1-5-1, Marunouchi, Chiyoda-ku, Tokyo 100-8405, Japan

<sup>4</sup> Brook Scientific Consulting LLC, Hilton Head Island, SC, USA



**Fig. 5** Identification of species-differences in  $\beta$ -lyase dependent formation of MFA and metabolites containing a  $\text{CH}_2\text{F}$  unit in cytosolic fractions of minipigs, rabbits and humans. Resonances were assigned to MFA ( $\delta = -217.1$  ppm;  $t, J_{\text{HF}} = 48.4$  Hz); unknown  $\text{CH}_2\text{F}$

unit containing metabolite (I) ( $\delta = -202.0$  ppm;  $t, J_{\text{HF}} = 48.5$  Hz); unknown  $\text{CH}_2\text{F}$  unit containing metabolite (II) ( $\delta = -208.2$  ppm,  $t, J_{\text{HF}} = 48.5$  Hz); unknown  $\text{CH}_2\text{F}$  unit containing metabolite (III) ( $\delta = -220.2$  ppm;  $t, J_{\text{HF}} = 48.5$  Hz)

holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.